



UNIVERSITÀ DI PARMA

ARCHIVIO DELLA RICERCA

University of Parma Research Repository

The transient gravitational-wave sky

This is the peer reviewed version of the following article:

Original

The transient gravitational-wave sky / Andersson, Nils; Baker, John; Belczynski, Krzysztof; Bernuzzi, Sebastiano; Berti, Emanuele; Cadonati, Laura; Cerdá Durán, Pablo; Clark, James; Favata, Marc; Finn, Lee Samuel; Fryer, Chris; Giacomazzo, Bruno; González, Jose Antonio; Hendry, Martin; Heng, Ik Siong; Hild, Stefan; Johnson McDaniel, Nathan; Kalmus, Peter; Klimentenko, Sergei; Kobayashi, Shiho; Kokkotas, Kostas; Laguna, Pablo; Lehner, Luis; Levin, Janna; Liebling, Steve; Macfadyen, Andrew; Mandel, Ilya; Marka, Szabolcs; Marka, Zsuzsa; Neilsen, David; O'Brien, Paul; Perna, Rosalba; Read, Jocelyn; Reisswig, Christian; Rodriguez, Carl; Ruffert, Max; Schnetter, Erik; Searle, Antony; Shawhan, Peter; Shoemaker, Deirdre; Soderberg, Alicia; Sperhake, Ulrich; Sutton, Patrick; Tanvir, Nial; Was, Michal; Whitcomb, Stan. - In: CLASSICAL AND QUANTUM GRAVITY. - ISSN 0264-9381. - 30:19(2013), p. 193002. [10.1088/0264-9381/30/19/193002]

Published

DOI:10.1088/0264-9381/30/19/193002

Terms of use:

openAccess

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

(Article begins on next page)

06 May 2023

The Transient Gravitational-Wave Sky

Nils Andersson,¹ John Baker,² Kris Belczynski,^{3,4} Sebastiano Bernuzzi,⁵ Emanuele Berti,^{6,7} Laura Cadonati,⁸ Pablo Cerdá-Durán,⁹ James Clark,⁸ Marc Favata,¹⁰ Lee Samuel Finn,¹¹ Chris Fryer,¹² Bruno Giacomazzo,¹³ Jose Antonio González,¹⁴ Martin Hendry,¹⁵ Ik Siong Heng,¹⁵ Stefan Hild,¹⁶ Nathan Johnson-McDaniel,⁵ Peter Kalmus,¹⁷ Sergei Klimenko,¹⁸ Shiho Kobayashi,¹⁹ Kostas Kokkotas,²⁰ Pablo Laguna,²¹ Luis Lehner,²² Janna Levin,²³ Steve Liebling,²⁴ Andrew MacFadyen,²⁵ Ilya Mandel,²⁶ Szabolcs Marka,²⁷ Zsuzsa Marka,²⁸ David Neilsen,²⁹ Paul O'Brien,³⁰ Rosalba Perna,¹³ Harald Pfeiffer,³¹ Jocelyn Read,³² Christian Reisswig,⁷ Carl Rodriguez,³³ Max Ruffert,³⁴ Erik Schnetter,^{22,35,36} Antony Searle,¹⁷ Peter Shawhan,³⁷ Deirdre Shoemaker,²¹ Alicia Soderberg,³⁸ Ulrich Sperhake,^{39,7,40} Patrick Sutton,⁴¹ Nial Tanvir,³⁰ Michal Was,⁴² and Stan Whitcomb¹⁷

¹*School of Mathematics, University of Southampton, Southampton, SO17 1BJ, UK*

²*Gravitational Physics Lab, NASA GSFC, Greenbelt, MD 20771, USA*

³*Astronomical Observatory, University of Warsaw, Al. Ujazdowskie 4, 00-478 Warsaw, Poland*

⁴*Center for Gravitational Wave Astronomy, University of Texas at Brownsville, Brownsville, TX 78520, USA*

⁵*Theoretical Physics Institute, University of Jena, 07743 Jena, Germany*

⁶*Department of Physics and Astronomy, The University of Mississippi, University, MS 38677, USA*

⁷*Theoretical Astrophysics 350-17, California Institute of Technology, Pasadena, CA 91125, USA*

⁸*Physics Department, University of Massachusetts, Amherst, MA 01003, USA*

⁹*Departamento de Astronomía y Astrofísica, Universidad de Valencia, Dr. Moliner 50, 46100 Burjassot, Spain*

¹⁰*Montclair State University, 1 Normal Ave, Montclair NJ 07043, USA*

¹¹*Department of Physics & Department of Astronomy & Astrophysics,*

The Pennsylvania State University, University Park, PA 16802, USA

¹²*CCS-2, Los Alamos National Laboratory, Los Alamos, NM 87545, USA*

¹³*JILA, University of Colorado and National Institute of Standards and Technology, Boulder, CO 80309, USA*

¹⁴*Instituto de Física y Matemáticas, Universidad Michoacana de San Nicolás de Hidalgo. Edificio C-3 Ciudad Universitaria 58040, Morelia, Michoacán, México*

¹⁵*SUPA, School of Physics and Astronomy, University of Glasgow, G12 8QQ, UK*

¹⁶*Institute for Gravitational Research, University of Glasgow, G12 8QQ, UK*

¹⁷*LIGO Laboratory, California Institute of Technology, Pasadena, CA 91125, USA*

¹⁸*University of Florida, PO Box 118440, Gainesville, Florida 32611, USA*

¹⁹*Astrophysics Research Institute, Liverpool John Moores University, Birkenhead, CH41 1LD, UK*

²⁰*Theoretical Astrophysics, IAAT, Eberhard Karls University of Tuebingen, Tuebingen 72076, Germany*

²¹*School of Physics, Georgia Institute of Technology, Atlanta, Georgia 30332, USA*

²²*Perimeter Institute for Theoretical Physics, Waterloo, Ontario, Canada*

²³*Department of Physics and Astronomy, Barnard College of Columbia University, New York, NY 10027, USA*

²⁴*Long Island University, Brookville, NY 11548, USA*

²⁵*Physics Department, New York University, New York, NY 10003, USA*

²⁶*School of Physics and Astronomy, University of Birmingham, Edgbaston, B15 2TT, UK*

²⁷*Department of Physics, Columbia University, New York, New York 10027, USA*

²⁸*Columbia Astrophysics Laboratory, Columbia University, New York, New York 10027, USA*

²⁹*Department of Physics & Astronomy, Brigham Young University, Provo, UT 84602, USA*

³⁰*Department of Physics & Astronomy, University of Leicester, University Road, Leicester, LE1 7RH, UK*

³¹*Canadian Institute for Theoretical Astrophysics,*

University of Toronto, Toronto, Ontario M5S 3H8, Canada

³²*Department of Physics, California State University Fullerton, Fullerton, CA 92831, USA*

³³*Department of Physics and Astronomy, Northwestern University, Evanston, IL 60208, USA*

³⁴*School of Mathematics and Maxwell Institute, King's Buildings,*

University of Edinburgh, Edinburgh EH16 5JN, Scotland, UK

³⁵*Department of Physics, University of Guelph, Guelph, Ontario, Canada*

³⁶*Center for Computation & Technology, Louisiana State University, Louisiana, USA*

³⁷*University of Maryland, College Park, MD 20742, USA*

³⁸*Harvard-Smithsonian Center for Astrophysics, 60 Garden st. Cambridge, MA 02138, USA*

³⁹*DAMTP, Center for Mathematical Sciences, Cambridge CB3 0WA, UK*

⁴⁰*CENTRA-IST, Lisbon, Portugal*

⁴¹*School of Physics and Astronomy, Cardiff University, Cardiff, CF24 3AA, UK*

⁴²*Albert-Einstein-Institut, Max-Planck-Institut für Gravitationsphysik, D-30167 Hannover, Germany*

(Dated: May 6, 2013)

Interferometric detectors will very soon give us an unprecedented view of the gravitational-wave sky, and in particular of the explosive and transient Universe. Now is the time to challenge our theoretical understanding of short-duration gravitational-wave signatures from cataclysmic events, their connection to more traditional electromagnetic and particle astrophysics, and the data analysis

techniques that will make the observations a reality. This paper summarizes the state of the art, future science opportunities, and current challenges in understanding gravitational-wave transients.

I. INTRODUCTION

The gravitational wave (GW) sky is an unexplored frontier which holds a great potential for discovery and a promise for understanding one of the most mysterious interactions of nature: gravity. Predicted by Einstein's theory of general relativity, GWs are ripples in the fabric of space-time, produced by the accelerated motion of masses. They carry information from the bulk, coherent motion of matter, complementary to the multi-wavelength electromagnetic spectrum of traditional astronomy and to the neutrinos and cosmic rays of particle astrophysics. Their observation will play a transformative role in our understanding of the Universe.

The existence of GWs was indirectly proven by over three decades of measurements of the orbit of the binary pulsar PSR1913+16, which has steadily been evolving due to the emission of gravitational radiation in agreement with the predictions of general relativity [1]. However, the direct measurement of GWs remains a challenge, due to their tiny amplitude once they reach Earth.

New and upgraded GW detectors are pursuing their first detection, which will transition gravitational physics to an observation-driven field and usher in a new GW astronomy. The instrumental landscape includes a new generation of ground-based laser interferometers [2–4], pulsar-timing arrays [5–7] and future detectors on the ground and in space [8–13]. In particular, the first generation of interferometric detectors has achieved design sensitivity [14–16], and next-generation ground-based interferometers are expected to be taking data within a few years.

The “perfect storm” in the transient sky of short-duration cataclysmic events is about to arrive, with GW observations from stellar core collapse, gamma-ray burst engines, rapidly rotating neutron stars (NSs) and mergers of compact object binaries. The storm will be fueled by GW observations from Advanced LIGO and Advanced Virgo, but the success in interpreting observations will hinge on our ability to model the complex physics at the heart of these transient astrophysical sources. This includes GW emission mechanisms that, in general, are not yet fully understood. The importance of this enterprise was highlighted in *New Worlds, New Horizons in Astronomy and Astrophysics*, a Decadal Survey of Astronomy and Astrophysics by the National Research Council [17].

This paper summarizes the current state of the art, future science opportunities and open challenges for GW transients. In §II, we review the main sources of the GW transient sky: mergers of compact objects, core collapse supernovae, and NS oscillations. For each burst source, we discuss its chances of detection and its connection with electromagnetic observations. In §III, we describe GW antennas as well as traditional astronomi-

cal observatories available today and in the near future. We elaborate on the detection challenges and open questions of the GW transient universe in §IV. Finally, §V summarizes our conclusions.

This review was conceived as the peroration of the *Gravitational Wave Bursts* workshop series. The first of these workshops was held in Chichen-Itza, Mexico on December 9-11, 2009 and the second in Tobermory, Scotland, on May 29-31, 2012. These meetings provided a forum for astrophysicists, GW data analysts and numerical relativists to explore transformative views of the GW transient sky, focused on: i) critical examination of current methodologies to model, detect and characterize transients, ii) current understanding of the physics behind burst sources, iii) requirements on detector technology and data analysis, and iv) imagining the future of GW transient science.

II. SOURCES OF GRAVITATIONAL WAVE TRANSIENTS

Compact objects such as NSs and black holes (BHs) will likely be protagonists in most of the astrophysical events detectable by the next generation of ground-based GW interferometers. Their role could start at birth (core collapse supernovae) or later in their life, either as members of binary systems or as isolated objects. This section provides an overview of what we know, observationally and theoretically, about compact objects as potential sources of transient GWs, including their predicted energetics and rates.

A. Compact Object Binaries and Short Gamma-ray Bursts

Binaries of coalescing compact objects are the main target of ground-based GW astronomy. In many instances, their gravitational waveform is expected to contain many cycles in the sensitive band of the detectors. Therefore, with the aid of well understood models (and assuming that Einstein's general relativity provides the correct description of gravity in dynamical, strong-field systems) they are ideal candidates for discovery via matched filtering. Given what we know about these sources and their merger rates deduced from known NS systems [18], binaries with either two NSs or a NS and a BH (also called *mixed binaries*) are expected to be bread-and-butter sources for ground-based detectors, even though recent population synthesis calculations suggest that BH-BH binaries may be more numerous than initially expected [19, 20]. Unlike binary BHs, binaries containing at least one NS have an additional appeal: their potential to act as the central engine of short

gamma-ray bursts (GRBs) (and possibly other observable electromagnetic signals). Short GRBs are bursts with duration shorter than ~ 2 seconds, with most of them lasting a few hundreds of milliseconds. The expected properties of a population of short GRBs from double compact object mergers have been estimated by several studies [21–23]. To date, only rough comparisons can be made between theory and observations [24, 25]. However, the theoretical predictions of compact object merger rates are in general consistent with the observed short GRB rates [23, 26].

There are several pieces of evidence supporting the association of short GRBs with the merger of double NS or mixed binaries [27]. One is that short GRBs do not seem to be associated with supernovae, and some of them explode in “dead” elliptical galaxies (i.e., galaxies with negligible ongoing star-formation). Long GRBs have distinctly different host galaxies from short GRBs, and they mainly occur in star-forming galaxies. Short GRBs appear more closely correlated with the rest-frame optical light (old stars) than the UV light (young massive stars). Furthermore, offsets of short GRBs relative to their host galaxy centers are significantly larger than for long GRBs [28].

The observations listed above are promising, but several challenges remain to establish an unequivocal association between short GRBs and compact binary mergers. As we will see below, the answer to these questions for now can only be obtained via sophisticated numerical models including the relevant microphysics. For example, a problem that can only be resolved via numerical methods is whether mergers can produce accretion disks massive enough to power the observed electromagnetic emission. Other demanding aspects of the bursts are precursors [29] and extended emission phases [30] that happen on timescales larger than 10 seconds: these timescales are beyond the reach of current simulations, which presently last less than a second. Looking forward, a smoking gun for the association between short GRBs and binary mergers will be the coincident detection of electromagnetic signals and GWs. Such a coincident detection is one of the most exciting multi-messenger observations that could occur in the advanced detector era.

1. Neutron Star-Neutron Star Binaries

The last few years have witnessed remarkable progress in fully general relativistic simulations of compact object binaries [31]. The first simulations of NS binaries were performed by Shibata and collaborators [32], but only recently have simulations been extended from the late inspiral up to the coalescence and eventual formation of a BH surrounded by a massive torus. Also recent is the inclusion of more sophisticated physics – realistic equation of state (EOS), magnetic fields and neutrino radiation – as well as the implementation of advanced numerical algorithms, such as adaptive mesh refinement

(AMR) techniques.

New physics and AMR have been essential ingredients to accurately predict GW emission and to establish a possible connection with short GRBs. In particular, having AMR in place in the numerical codes has allowed various groups to model the inspiral and merger of NSs [34–37] and to study the formation of hydrodynamic instabilities in the presence of magnetic fields [34, 38, 39]. Similarly, the use of a realistic EOS has led to the suggestion [40, 41] that there are two broad classes of NS binary mergers. Binaries with initial total masses above $\sim 2.8 - 3.2M_\odot$ (depending on the EOS) promptly form a BH soon after the merger. On the other hand, binaries with lower masses yield a metastable, hypermassive NS before collapsing to a BH. This difference has very important observational consequences. In the case of massive binaries, the GW signal shows a quick transition from “chirping” during the inspiral to the characteristic quasinormal ringdown of the final BH [42]. On the other hand, if the merger yields a hypermassive NS, the chirp in the GWs is followed by quasi-periodic oscillations at frequencies $2 - 4$ kHz: see e.g. Figure 1, adapted from [33]. Unfortunately this quasi-periodic signal will only be detectable by Advanced LIGO and Virgo interferometers if the merger takes place within ~ 20 Mpc [43, 44], but these systems could be an interesting target for future detectors.

Recent studies [45, 46] have also addressed the association of binary NS mergers with short GRBs in the more general case of unequal-mass binaries. These investigations showed that the formation of a massive torus, which could lead to the emission of gamma rays, leaves a characteristic signature in the GW spectrum, consisting of an exponential decay followed by a hump in the $\sim 2 - 7$ kHz range. The cut-off frequency and the frequency and amplitude of the hump can be directly associated to the NS EOS and to the mass of the disk. While Advanced LIGO and Virgo may not be sufficiently sensitive at these frequencies, a third-generation detector (such as the Einstein Telescope) would be sensitive to sources within 100 Mpc. Moreover, simulations have shown that unequal-mass binaries can produce tori with masses up to $\sim 0.35M_\odot$, which would be more than sufficient to power short GRBs [33]. Notice however that simple equipartition arguments would suggest that even tori with masses as low as $\sim 0.01M_\odot$ may be sufficient to power short GRBs (see e.g. [47] for a discussion).

Present estimates suggest that observations of lowest-order tidal perturbations in NS mergers with third-generation detectors would yield measurements of NS radii with ~ 1 km precision [48, 49], while observations of strong-field tidal effects during mergers might allow similar accuracies even with second-generation detectors if sufficiently accurate models are available [50]. Such GW observations, which probe the bulk properties of neutron stars and could make it possible to constrain the EOS of matter in the NS core, will be complementary to electromagnetic observations [51, 52], which are sen-




FIG. 1. GW signal from the merger of equal-mass binary NSs. The left panel refers to a “high-mass” case with a total gravitational mass of $3.23M_{\odot}$; the right panel, to a “low-mass” binary with a gravitational mass of $2.69M_{\odot}$. In the “high-mass” case the total mass of the system is large enough to produce a prompt collapse to BH soon after the merger, and the GW signal is characterized only by inspiral, merger and BH ringdown. In the right panel, on the other hand, the mass of the system is lower; a hypermassive NS is formed and survives for ~ 120 ms before collapsing to a BH (note the very different time scale between the two panels). In the right panel, different insets show zoom-ins on different parts of the signal. The bottom-left inset shows the inspiral and merger on the same time scale as the left panel. The top inset shows the signal emitted by the newborn hypermassive NS, which has a peak frequency at ~ 2.5 kHz. The bottom-right inset shows the final collapse to a BH. The figure was produced using data from [33].

sitive to surface properties. This enterprise will require a large number of GW templates. To reduce the computational burden, there have been successful attempts to construct semi-analytical GW templates (e.g. using the effective one-body approach) that include the influence of tidal deformation [53–55]. More recent studies in the conformally-flat approximation [56, 57] have suggested that the GW signal emitted by the hypermassive NS formed after merger may also be used to constrain the NS EOS. If the result of the merger is a long-lived hypermassive NS, the GW signal may be used to measure the NS radius with an accuracy of up to ~ 200 m already with Advanced LIGO [56, 57]. While this is a very interesting result, a more accurate treatment of general relativistic effects and magnetic fields may affect the evolution of the hypermassive NS [34, 58] and make such measurements more difficult.

Our understanding of the potential role of magnetic fields in the dynamics of NS binaries has significantly improved in recent times [38, 39, 58–62]. These studies have provided additional support to the view that the merger of magnetized NSs can provide the central engine for short GRBs; furthermore, one could have potentially observable emissions even in cases where a SGRB is not realized [60–62]. Unfortunately, the studies also showed that the effects of magnetic fields on the GW signal are

most appreciable at frequencies larger than ~ 1 kHz, where Advanced LIGO and Virgo are less sensitive.

Simulations of NS mergers that include the effects of neutrino cooling are in their infancy. Pioneering work in this direction [43, 44, 63] shows that for a stiff, finite-temperature EOS and with neutrino cooling, a hypermassive NS is the canonical outcome of the merger of NS binaries with masses smaller than $3.2M_{\odot}$, and that thermal pressure support may be important. The neutrino luminosity associated with these events could potentially be detected for mergers within 5 Mpc by hyper-Kamiokande [44]. Finally, studies of the GW emission from NS mergers on eccentric orbits have also begun [64, 65]. These systems may be formed via dynamical capture in dense stellar environments, and they may account for a fraction of NS binaries [66].

2. Neutron Star-Black Hole Binaries

In the last few years, the interest in modeling mixed binary systems comprised of a BH and a NS has also intensified (see [67] for a recent review). The first simulations tracked the merger of a NS with a non-spinning BH [68–70]. These early simulations showed that the merger produced a torus with mass $\sim 0.2M_{\odot}$ when the

BH and the NS have comparable masses. Subsequent studies with higher numerical accuracy excluded the possibility of short GRBs in mixed binary mergers if the BH is non-spinning [36, 71–73], even in the case of equal-mass systems [72].

If the BH is spinning, the merger can potentially power a short GRB. Simulations show that a disk with mass $\sim 0.2M_\odot$ can be formed for binary mass ratios of 1/3 and BH spins of $a/M_h = 0.75$ [74], where $a = J/M$ is the Kerr spin parameter (in geometrical units $G = c = 1$). Recently, the case of NSs merging with $10M_\odot$ BHs was explored in [75]. If the BH is rapidly rotating ($a/M_h \sim 0.7 - 0.9$) the merger can result in accretion disks massive enough to power a short GRB [75]; however, we remark once again that even less massive tori may be able to power short GRBs in the presence of instabilities [47]. Furthermore, neutron-rich ejecta from BH-NS systems are possible, and it has been shown that the calculated flux and the time to return to the central engine are consistent with models for extended emissions from the r-process [76].

Analytic models have also been developed to compute the mass of the disk that can be formed after merger [77, 78], as well as other features of the merger remnant [79]. These studies have the main advantage that they allow us to explore a larger portion of the parameter space. They confirmed that massive tori can be formed even at low mass ratios ($q \sim 0.1$) if the BH is rapidly spinning [77, 78], and therefore that these binaries could in principle power short GRBs.

GWs emitted from mixed binary systems can be grouped in three broad classes depending on their behavior near the innermost stable circular orbit (ISCO) [80]:

- Type I: the NS is disrupted outside the ISCO.
- Type II: the mass transfer from the NS to the BH takes place close to the ISCO.
- Type III: the NS is not disrupted outside the ISCO, and all of the matter falls immediately into the BH.

While gravitational waveforms from type-III mergers are difficult to distinguish from BH binary systems, type I and II will be sufficiently distinct that they could provide information about the mass ratio and the NS compactness. The likely frequency of these distinctive signatures is above 2 kHz, i.e. within the reach of third-generation detectors, such as the Einstein Telescope. In the case of non-spinning BHs and for large mass ratios between ~ 0.3 and 1 (i.e., type-I/type-II mergers), tidal deformations induced in the NS during the inspiral may also allow for Advanced LIGO measurements of the NS radius (and hence for constraints on the EOS) with accuracy $\sim 10\% - 50\%$ for sources located at a distance of 100 Mpc [81–83].

While early simulations of mixed binary mergers adopted a simple ideal-fluid EOS, recent work accounts for more realistic EOSs [84–86], the effects of the orientation of the BH spin on the formation of the torus

and GW signatures [87], and the presence of magnetic fields [88, 89]. Magnetic fields do not seem to have a detectable GW signature in the case of mixed binaries, but they could provide a mechanism for the production of relativistic jets when a torus is formed after the merger, and various models predict that they could induce possible observable electromagnetic counterparts [90–93].

Finally, recent studies considered eccentric orbits in mixed binary mergers [94, 95]. In particular, these simulations addressed the effect of eccentricity on the formation of the massive torus and on GW emission. While binaries in quasi-circular orbits emit a periodic signal during their inspiral phase, eccentric binaries emit a series of quasi-periodic GW bursts, detectable by Advanced LIGO and Virgo up to distances of 300 Mpc.

3. Event Rates

Techniques for estimating rates of compact binary mergers have been recently summarized in [96]. The rate of NS-NS binary mergers in our Galaxy can be estimated from measured parameters of known binary pulsars [97]. Given the lack of direct observations of NS-BH systems, their rates must be estimated by modeling, typically involving population-synthesis studies of large numbers of simulated binaries [98], or attempts to predict the future evolution of specific observed systems that represent possible progenitors of compact binaries [99].

Unfortunately, both methods currently suffer from significant uncertainties. Population-synthesis results depend critically on assumptions about common-envelope evolution, typical supernova kicks, mass-loss rates, metallicity, and other astrophysical parameters [19, 100]. Extrapolations from binary pulsar observations have fewer free parameters, but their accuracy is limited by small-number statistics due to the paucity of observed binary NS systems in the Galaxy, the imperfect understanding of selection effects in pulsar surveys, and uncertain knowledge of the pulsar luminosity function.

According to the compilation [101], NS-NS merger rates plausibly range from 1 to 1000 mergers per Milky Way Galaxy per million years. This range is also consistent with extrapolations from the observed rate of short GRBs assuming a relatively high correction for beaming [26]. Meanwhile, NS-BH merger rates fall in the range 0.05 to 100 per million years in the Galaxy.

The conversion of merger rates to detection rates depends on the assumed detector sensitivities, data quality, and details of the search pipelines. The uncertainties in these factors are typically small relative to the uncertainties considered above, but additional astrophysical uncertainties encountered when scaling up from the Galaxy, such as including the contribution of low-metallicity environments or elliptical galaxies, could be more significant [102]. The merger rate ranges quoted above correspond to detection rates of 0.04 to 400 per year for NS-NS binaries and 0.2 to 300 per year for NS-BH binaries in the

era of advanced GW detectors operating at full sensitivity [101].

B. Binary Black Hole Mergers

The defining characteristic of BHs is their event horizon: a “surface of no return,” from within which not even light can escape. Until now, we have been able to infer the existence of BHs only indirectly, in particular by modeling phenomena associated with the neighborhood of the putative BH horizons [103]. The evidence gathered so far is from the behavior of astrophysical objects, matter or fields that cannot be explained by other means than by appealing to the presence of a BH, under the assumption that Einstein’s theory of gravity is correct. Examples are the emission from active galactic nuclei, the orbits of stars at the center of our Galaxy, X-ray sources and tidal disruptions of stars, to name a few. One rather extreme point of view is that no electromagnetic observation will ever provide conclusive proof of the existence of BHs [104]. On the other hand, there is general consensus that the detection and characterization of GWs from the merger of two BHs will offer compelling evidence for their existence: see e.g. [42, 105] and references therein.

Binary stellar mass BHs can be formed either through the evolution of isolated binaries in galactic fields, or through dynamical formation scenarios in dense stellar environments, such as globular clusters or galactic nuclear clusters (see [96, 101] and references therein). Due to the lack of direct observations of any binary BHs, predictions about rates and mass distributions of these systems must rely on simulations.

In Section II A 3 we referred to population-synthesis models for the evolution of isolated compact-object binaries. These models have particularly large uncertainties in the case of binary BHs, including the effects of metallicity (lower metallicity tends to decrease mass loss through stellar winds and increase the number of merging binary BHs), the uncertainty in supernova birth kicks for BHs (higher kicks may disrupt binaries), and the uncertain future of binaries that enter the common envelope as they are going through the Hertzsprung gap (such binaries may merge directly, without a GW signature). According to the compilation of predictions in [101], advanced detectors may observe GWs from merging binary BHs at a rate between one detection in a few years and a thousand detections per year. More recent simulations considering a wider range of updated models are presented in [100]. These simulations indicate that rate uncertainties still span several orders of magnitude, but also that rates appear more promising than in the past. Attempts to model the future evolution of BH–Wolf-Rayet binaries IC 10 X-1 and NGC 300 X-1 also indicate that advanced detectors may observe hundreds of events per year [106], even though the precise modeling of the evolution of such systems is still a difficult task.

Another possible channel to produce observable binary

BH mergers consists of dynamical interactions in globular clusters and nuclear star clusters. In these systems, BHs are likely to sink to the center through mass segregation and replace other members of existing binary systems via three-body encounters, leading eventually to binary BH mergers. Several simulations (see e.g. [107–110]) have indicated that, although many BHs could be ejected from globular clusters during three-body encounters, dynamically formed BH-BH binaries could make significant contributions to the overall rates of detected systems.

Most BH binaries produced in population synthesis models or via three-body encounters in globular clusters and nuclear star clusters have masses such that the binaries will inspiral in the band of interest for Earth-based GW detectors; some of them may also produce detectable ringdown signals in band. Post-Newtonian approximations and progress in numerical relativity since the 2005-2006 breakthroughs [111–113] are supplying accurate knowledge of the type of signal emitted in the coalescence, by providing waveforms with enough cycles to cover the inspiral, merger and ringdown (see e.g. [105, 114, 115] for recent reviews). Knowledge of the waveform allows for matched-filtering searches for these signatures in GW data. Efforts to create analytical and/or phenomenological models calibrated to numerical-relativity data are ongoing: see e.g. [116–121].

Beyond the mass range of stellar-mass BHs, binaries involving intermediate-mass black holes (IMBHs) could represent exciting GW sources. Detectable binaries in advanced LIGO and Virgo would have total mass in the range between ~ 100 and ~ 500 solar masses. Both theoretical formation scenarios and observational evidence for IMBHs are topics of active research and debate [122, 123]. If IMBHs have a non-negligible occupation number in globular clusters, they could capture stellar-mass NSs or BHs, with GWs from the ensuing intermediate mass ratio inspirals being detectable at rates of up to tens per year [124, 125]. It may also be possible to detect mergers of two IMBHs [126, 127]. The highly uncertain rates of these processes are summarized in [101].

The characteristic chirp-like signal from binaries in a quasi-circular inspiral could be radically modified if the BHs merge in a highly eccentric, precessing orbit. Such orbits could be the result of scattering events expected in the environment of dense galactic cores. The resulting GWs will appear as bursts of radiation near periastron, followed by quiescent phases while the BHs travel to and return from apoastron. The time elapsed between subsequent bursts decreases as the binary hardens and the eccentricity decreases [128–134]. Depending on the masses of the BHs, GWs from highly eccentric binaries should be visible by both ground- and space-based interferometers [135, 136].

C. Core-Collapse Supernovae and Long Gamma-ray Bursts

Core collapse supernovae (CCSNe) and long-duration gamma-ray bursts (LGRBs) share a common origin: the collapse of stars with masses $\gtrsim 8 M_{\odot}$. Observational evidence comes from the positional and temporal association of several nearby LGRBs with supernovae of Type Ic: see [137, 138] for reviews. These SNe associated with GRBs are distinguished from other core-collapse events based on (i) the absence of hydrogen and helium in their optical spectra [139], characterized predominantly by broad absorption features of intermediate mass elements, and (ii) a strong non-thermal afterglow component best studied at radio wavelengths [140]. Long-term monitoring of the afterglow reveals that the LGRB is characterized by a bi-polar relativistic outflow, and jet opening angles $\lesssim 30^{\circ}$ are commonly inferred [141]. Intriguingly, comparing the collimation-corrected rates of LGRBs and Type Ic SNe, we find that *most* LGRBs are associated with a SN, but less than 1% of SN Ic are associated with a LGRB. This is confirmed through detailed radio studies of local Type Ic SNe, which indicate that only $\sim 0.7\%$ of SNe Ic drive relativistic outflows, some of which do not give rise to detectable ($E_{\gamma} \gtrsim 10^{48}$ erg; 25-150 keV) gamma-ray emission [142, 143].

While the progenitors of SNe Ic and/or LGRBs have yet to be directly detected in pre-explosion imaging [144], theoretical considerations point to massive stars that have been stripped of their hydrogen envelope prior to explosion, either by their own strong radiation-driven stellar winds [145] or through the interaction with a close binary companion [146]. The critical ingredient that enables only 1% of SNe to produce GRBs remains unclear, but a key component is probably low metallicity ($Z \lesssim 0.5 Z_{\odot}$), allowing the stellar core to retain angular momentum by suppressing the line-driven winds [147]. With the advent of new wide-field surveys (e.g., Pan-STARRS, Palomar Transient Factory), the rate of CCSN discoveries in metal-poor galaxies is growing. Combined with radio follow-up observations, the metallicity dependence of relativistic outflows in CCSNe can be directly tested.

GRB-associated SNe are not the only explosions with evidence for asphericity. Late-time spectroscopy of CC-SNe in the nebular phase [148], polarization measurements of local CCSNe [149], and detailed high spatial resolution studies of Galactic SN remnants such as Cas A [150] all suggest that ejecta asymmetries are in fact common place. Thus, it remains unclear what distinguishes the progenitors of ordinary CCSNe from GRBs, and it could be that a significant fraction of CCSNe (not just those associated with *detected* gamma-ray bursts) could experience exotic explosion mechanisms.

1. Theoretical Modeling

At the end stage of stellar evolution, the core of a massive star is supported against gravity by the pressure of relativistically degenerate electrons. Collapse is initiated when the core exceeds its effective Chandrasekhar mass and continues until the inner core reaches nuclear density. There, the nuclear EOS stiffens, leading to core bounce and the formation of the hydrodynamic bounce shock. The shock runs into the supersonically collapsing outer core, losing its energy to the break-up of infalling heavy nuclei into nucleons and to neutrinos that are made by electron capture in the region behind the shock, and stream out freely as the shock reaches regions of low neutrino optical depth. The shock stalls, turns into an accretion shock, and must be revived by the *core-collapse supernova mechanism* to drive a core-collapse supernova explosion. This is the basic picture that has been established since Bethe's authoritative 1990 review [151].

A variety of mechanisms have been proposed in the literature (see, e.g., [152] for a recent review) and all leading candidates involve GW-emitting aspherical dynamics. The *neutrino mechanism* relies on the net deposition of energy by charged-current neutrino absorption in the region immediately behind the stalled shock. While the neutrino mechanism fails to blow up ordinary massive stars in spherical symmetry, the multi-dimensional phenomena of convection, turbulence, and the standing-accretion shock instability (SASI, an advective-acoustic instability of the stalled shock) very likely enhance the neutrino mechanism's efficacy [153–159].

The *magnetorotational mechanism* relies on rapid rotation and magnetic field amplification due to flux compression in collapse, rotational winding, and the magnetorotational energy that converts free energy of differential rotation into magnetic field [160–164]. These processes may lead to magnetic fields with strengths of order 10^{15} G, which would be sufficient to launch bipolar magnetohydrodynamic jets, leading to an energetic, strongly aspherical explosion [165]. The key ingredient required for the magnetorotational mechanism is rapid progenitor star rotation, but most massive stars, perhaps up to 99%, are presently expected to be slow rotators [166, 167].

A third potential way of driving core-collapse supernova explosions is the *acoustic mechanism* proposed by [168, 169]. In their simulations, SASI-modulated turbulence and accretion downstreams hitting the protoneutron star (PNS) excited pulsations (predominantly of $l = 1$ spatial character) of the latter, that grew to non-linear amplitudes and dissipated in sound waves. Propagating along the density gradient behind the stalled shock, the sound waves steepened to secondary shocks, injecting additional heat into the postshock region, eventually leading to explosion. This mechanism is robust, but there are many unresolved issues with it. Explosions occur quite late and would imply NS masses and nucleosynthetic yields that are likely inconsistent with observations. Only one group and code have produced

this mechanism to date. Though many see the necessary ingredients for this mechanism, including excitation of PNS pulsational modes, it is not clear whether the amplitudes obtained by [168, 169] are produced in nature. Furthermore, [170] showed that nonlinear parametric instabilities may limit the oscillation amplitudes of the PNS by funneling oscillation power into daughter modes.

The details of the long GRB central engine may be as uncertain as the core-collapse supernova mechanism, but the relativistic beamed outflows observed from GRBs strongly suggest that rapid rotation plays a major role in the central engine.

In the collapsar scenario, outlined first by [171], a rotating core-collapse supernova fails to explode or explodes weakly or very aspherically, leading to BH formation before (type-I collapsar) or after (type-II collapsar) an explosion by fallback accretion. Eventually, typically seconds after BH formation [172], an accretion disk is expected to form near the BH. Accretion energy or extracted BH spin energy, mediated via MHD processes [173, 174] and/or neutrino pair annihilation [175, 176], may then drive the relativistic GRB outflow, while MHD disk winds and viscous heating may power a GRB-accompanying energetic core-collapse supernova explosion [177, 178].

The rapid progenitor rotation required in the collapsar scenario may lead to an energetic magnetorotational explosion, preventing BH formation [179]. This possibility gives rise to the competing millisecond protomagnetar model for the long GRB central engine [180–182]. In this model, a magnetorotational core-collapse supernova explosion excavates the polar regions, allowing the driving of an ultra-relativistic wind by the spin-down of the strongly magnetized neutrino cooling protoneutron star (the protomagnetar). The protomagnetar model is able to explain the prompt GRB emission, and prolonged magnetar activity may explain long-duration X-ray afterglow observed in long GRBs.

2. Gravitational Wave Emission

The ubiquitous aspherical dynamics in stellar collapse, core-collapse supernovae and long GRBs gives rise to bursts of GWs with typical durations from milliseconds to seconds, whose waveforms are impossible to predict precisely by simulations. The reason is that much of the GW emission is influenced or dominated by stochastic dynamics (i.e. turbulence), and also that much of the input physics (e.g., the nuclear EOS) and the initial conditions are complicated and impossible to know exactly.

Aspherical stellar collapse was early on considered as a source of detectable GWs [183, 184] and has been studied extensively: see [185–187] for recent reviews. In nonrotating or only slowly rotating core-collapse supernovae, the GW emission is dominated by convective overturn in the PNS and in the region behind the shock, modulated by the SASI and enhanced by fast accretion downstreams

that are decelerated in the stably stratified outer layers of the PNS. The emitted GW signal has random polarization, a broad spectrum with power at $\sim 100 - 1000$ Hz, and dimensionless strain amplitudes of order 10^{-22} at a source distance of 10 kpc [155, 188–191]. Contributions at low frequencies ($\lesssim 30$ Hz) come from anisotropic emission of neutrinos [188, 190–192] and explosion asphericities [155, 189, 190].

If the neutrino mechanism lacks efficacy and the explosion is delayed to late time, the strong PNS oscillations associated with the acoustic mechanism may be excited. Their quadrupole components emit GWs at momentarily fixed (secularly changing due to changes in the PNS structure) frequencies of $\sim 600 - 1000$ Hz, with strain amplitudes of $10^{-21} - 10^{-20}$ at 10 kpc [186, 193]. So far, all simulations of these pulsations have been axisymmetric, predicting linearly polarized signals, but in 3D correlated emission in the second GW polarization can be expected.

Rapid rotation, if present, will lead to a characteristic burst of GWs emitted at core bounce, when the inner core undergoes the greatest acceleration. This signal has been shown to be linearly polarized (i.e., the dynamics is axisymmetric) and increases in amplitude with increasing initial inner core angular velocity, up to the point where centrifugal forces become dominant and decelerate the bounce dynamics [194–198]. Typical signal amplitudes are of order 10^{-21} at 10 kpc for cores with precollapse central spin periods of 2 – 4 s; the GW emission peaks around 700 – 800 Hz, decreasing to below ~ 200 Hz for very rapid rotation. Simulations that take into account magnetic fields found that extreme precollapse iron core fields in excess of 10^{12} G would be necessary to modify the bounce dynamics and GW signal [164, 199–202]. More moderate initial fields can be amplified in the postbounce phase and will modify the postbounce dynamics and GW signal.

Subsequent to core bounce, nonaxisymmetric rotational instabilities may develop. These require rapid spin and/or differential rotation. A ratio of rotational kinetic to gravitational energy $T/|W|$ above $\sim 27\%$ is required for a classical high- $T/|W|$ dynamical instability, that leads to an $m = 2$ deformation of the PNS. A secular instability (driven by GW radiation reaction or viscosity) may set in at $T/|W| \gtrsim 14\%$ [203]. Typical rapidly spinning cores lead to PNS with $T/|W| \lesssim 10\%$ [194]. Core collapse naturally produces a nearly uniformly spinning PNS core with a strongly differentially rotating outer mantle [167]. This differential rotation can drive a rotational shear instability leading to angular momentum redistribution, nonaxisymmetric deformation and GW emission [197, 198, 204–210]. Typical signal characteristics are strain amplitudes of order 10^{-21} at 10 kpc and quasi-periodic emission at twice the frequency of the unstable mode – typically $\sim 800 - 1000$ Hz [197, 198, 209] – for a duration of 10 – few 100 ms.

After the onset of an explosion, the GW signal emitted by dynamics between the PNS core and the shock will subside quickly, leaving the more gradually decaying

GW signal from PNS convection and, potentially, nonaxisymmetric rotational dynamics behind. If the explosion fails, a BH forms after $\sim 1 - 3$ s (the exact time is determined by the nuclear EOS and the progenitor structure). If the PNS is spinning, this will give rise to a second pronounced peak in the GW signal, with strain of order 10^{-20} at 10 kpc and most GW power at frequencies above $1 - 2$ KHz [211].

In a collapsar-type long GRB, the GW emission will be very similar to a rapidly spinning core-collapse supernova up to BH formation. The latter will be followed by a multi-second GW-silent phase after which instabilities in the inner accretion disk and/or outer accretion torus may give rise to GW emission lasting, possibly, for the duration of the GRB [212–214]. Detailed waveforms of such instabilities have yet to be predicted by simulations.

In a millisecond-protomagnetar long GRB, the signal from BH formation and the GW-silent phase would be absent. The GW emission due to nonaxisymmetric rotational dynamics of the protomagnetar may continue for the duration of the GRB and, if the instability is secular, possibly throughout the early afterglow phase [215, 216], and may be detectable by advanced LIGO out to ~ 100 Mpc [215].

D. Isolated Neutron Stars

The NS menagerie offers a rich variety of electromagnetic phenomenology and possibilities for GWs. In addition to the binary NS coalescences discussed in Section II A, the birth of a NS following the death throes of a medium sized star in a supernova explosion may also be visible with advanced detectors. Each scenario may (depending on the maximum NS mass) lead to a hot, possibly rapidly spinning, remnant with violent dynamics. Initially this remnant is opaque to neutrinos, but after a few tens of seconds [217–219] it becomes transparent and cools, the thermal pressure drops and a NS is formed. During the initial phase, the GW signature of the new-born NS may evolve considerably [220], due to changes in thermal gradients and interior composition. This evolution, in fact, continues for the first few months of the NS life, as the crust freezes and the various superfluid/superconducting components establish themselves. At the end of the process, a mature NS has a complex structure, the modeling of which requires an understanding of much extreme physics.

In this section, we will focus on GW burst signals from isolated NSs. We take the, possibly simplistic, view that the related phenomena can be understood in terms of the star’s oscillation modes. This is certainly the case for many of the mechanisms that have been discussed in the literature, ranging from various mode-instabilities to magnetar flares and radio pulsar glitches. The interest in instabilities is natural, since they provide an explanation for the excitation of the modes and the associated GW signal. Similarly, it is reasonable to consider scenarios

associated with known electromagnetic phenomena, like magnetar flares and pulsar glitches. There are, however, issues with each of these scenarios. In the first case, the presence of an instability will not guarantee a detectable GW signal. The relevance of the mechanism depends on the physics that counteracts an unstable, growing mode. This involves both the mechanics of the problem – whether non-linear hydrodynamics saturates the instability [221–224] – and microphysics, as encoded in the relevant viscous damping channels [225]. While we have made good progress on understanding such issues in the last few years, it is clear that many challenges remain. In the second case, most estimates are based on simple energetics. However, the link between the observed electromagnetic signal and any GWs that may be generated by the underlying mechanism is not at all clear. This is rather obvious, since the detailed mechanisms leading to observed flares and glitches remain rather poorly modelled. Most current estimates are based on plausibility arguments. To make progress we need a better understanding of these enigmatic events. That this is a challenge is clear from the fact that the pulsar glitch mechanism and high-energy emissions from pulsars are not well understood, despite four decades of study [226].

NSs may radiate GWs through a range of mechanisms. For traditional reasons, the associated signals tend to be categorized either as “bursts” or “periodic” signals (mainly because of the different data analysis strategies used to look for the signals). However, this division is rather arbitrary. For example, rotating NSs with quadrupolar deformations (linked to the geological history of the elastic crust or the structure of the magnetic field) would produce continuous quasi-sinusoidal GW signals. Such signals have already been the subject of GW searches [227–229]. However, it is by no means clear that the signal will remain unchanged over a long term observation lasting months to years. In fact, one may argue that NS “mountains” ought to be transient, evolving due to plastic flow [230]. However, as the timescale and detailed behavior of the evolution is essentially unknown, this possibility has not been considered so far. A closely related problem concerns NSs that interact with their environment, as in the case of accretion from a binary partner in a Low-Mass X-ray Binary. While it is natural to assume that the accretion of material leads to some level of quadrupole deformation and GW emission [231], it is far from clear how such “mountains” are established and to what extent they evolve as the accretion rate changes. This is a key question that needs to be addressed if we are to search for signals from such systems. The problem is also linked to that of instabilities evolving on a secular timescale, as in the case of f- and r-modes discussed below. Instabilities may trigger a violent behaviour, but they may also be rather subtle, leading to the system simmering on the threshold of stability. The latter behaviour is, in fact, what is expected for the instability associated with inertial r-modes [222, 223, 232].

While searches for GWs from NSs have already been

performed on data from first-generation detectors [227–229, 233–237], it is generally expected that such signals will require third-generation detectors like the Einstein Telescope. A review of relevant sources of GWs, including isolated NSs, for the Einstein Telescope can be found in the design study [238] as well as recent review articles [239, 240]. Search techniques for long-duration transients have been proposed in [e.g., 241, 242]. Here, we will present a brief overview of the main mechanisms for “burst” emission from isolated NSs touching on the science returns for observing such signals, and highlighting issues that require further attention.

1. *Instabilities*

NSs may suffer various instabilities as they evolve from hot remnants to cold mature objects. Some of these instabilities may be efficient GW emitters and so are of obvious interest for GW astronomy. The relevant instabilities can be broadly divided into two classes. Dynamical instabilities tend to grow rapidly, and do not require additional “physics” for their existence. The most commonly considered such instability is the bar-mode instability associated with the star’s fundamental (f-) mode. Secular instabilities, on the other hand, are relatively subtle. They owe their existence to dissipative mechanisms and tend to grow on the associated dissipation timescale. The most important such instabilities (in the present context) are driven by GW emission via the so-called Chandrasekhar-Friedman-Schutz mechanism [243, 244] and are associated with the f-mode and the inertial r-mode in a rotating star. The basic picture is that an instability sets in at some threshold, say above a critical rotation rate, leading to the growth of a non-axisymmetric perturbation. The existence of an instability and the early phase of its evolution are relatively easy to establish, since they can be studied within linear perturbation theory. Once the growing mode reaches a sizeable amplitude the situation becomes less clear, as one must account for the full nonlinear dynamics. At some point one would expect the instability to saturate. Understanding the mechanism for, and level of, saturation is key if we want reliable estimates of the emerging GWs.

The dynamical bar-mode instability has been studied in some detail via numerical simulations. As anticipated in Section II C 2, this instability sets in once the ratio of rotational kinetic energy (T) to gravitational binding energy ($|W|$) exceeds a certain threshold. At that point, the f-mode grows and deforms the star into a (rotating) bar-shape. This would be a very efficient configuration for emitting GWs. However, it is not clear that real NSs exhibit this instability. The main problem is that the critical threshold requires a significant amount of differential rotation. Uniformly rotating NSs can not get near the critical value of $T/|W|$, because they reach the break-up limit before this happens. There are, of course, situ-

ations where differential rotation is expected to develop, most notably a few milliseconds after core-bounce in a supernova or tenths of seconds later, as the NS contracts due to cooling by neutrino emission. Present simulations [245–247] suggest that this may lead to $T/|W|$ becoming large enough to trigger the bar-mode instability, but unfortunately the system evolves away from this regime rather quickly. Recent simulations also cast doubt on the notion that the unstable bar-mode would last for many rotations [246], as required to make the effective GW amplitude detectable from sources outside our galaxy. A closely related, somewhat more subtle instability may be more important. There is evidence that instabilities may be triggered at much lower values of $T/|W|$, provided that the system exhibits significant differential rotation [248–252]. This class of instabilities is much less well understood at the moment. Most importantly, we need to establish whether real astrophysical systems may evolve into the relevant part of parameter space.

In the last few years, the main focus has been on secular (GW driven) instabilities. This kind of instability sets in when the pattern speed of a given modes changes from counter- to co-rotating with respect to an inertial frame. This effectively means that the system radiates positive angular momentum, drawn from a negative angular momentum reservoir, leading to a runaway process.

Early work on this mechanism focused on the instability of the f-mode in Newtonian NS models [253–256]. The results suggest (perhaps somewhat optimistically) that the unstable modes could be observable from sources beyond our Galaxy. However, the f-mode instability only operates near the mass shedding limit, so NSs would have to be born rapidly spinning for the mechanism to kick in during their early life. Current observations suggest that the subset of NSs born spinning sufficiently fast may be rather small [257], but this is certainly not well understood at the present time. Another problem for the f-mode instability is that it may be completely quenched by dissipation. In particular, the so-called mutual friction associated with superfluid vortices may suppress the instability once the star cools below the threshold for superfluidity [255, 258]. This means that the f-mode instability is unlikely to operate in mature NSs, e.g. ones spun up by accretion in a Low-Mass X-ray Binary. Our understanding of the f-mode instability has improved significantly recently, with accurate numerical simulations both at the linear and nonlinear level. In particular, we now have a clearer picture of the instability for relativistic stars [259, 260]. These results indicate an enhancement of the f-mode instability due to relativistic effects, renewing interest in the mechanism as a source for gravitational radiation [261].

In the last decade most work on secular instabilities has focused on the inertial r-modes. These modes are interesting as they radiate mainly through current multipoles, not mass multipoles (as in the case of virtually all other GW sources). That these modes would also be unstable due to the emission of GWs came as some sur-

prise, and by now many aspects of the associated instability have been considered: see [225, 262] for exhaustive reviews. The r-mode instability is interesting for many reasons. It may provide a natural explanation for the absence of NSs spinning faster than 720 Hz by preventing further spin-up once an accreting (recycled) NS reaches the instability threshold. This mechanism would lead to NSs in Low-Mass X-ray Binaries being interesting targets for GW searches [263]. However, due to the varying accretion rate in these systems and the many unknown parameters, such searches will be very difficult [264]. As in the case of the f-modes, the unstable r-modes are counteracted by a range of dissipative mechanisms. It is generally thought that the most important damping mechanisms are associated with i) a viscous boundary layer at the crust-core interface, ii) superfluid mutual friction, and iii) hyperon bulk viscosity in the deep core of the star. The star’s magnetic field may also have a decisive importance; this issue has not been studied in sufficient detail yet, but see [265]. Very recent work [266], comparing the predicted r-mode instability window to observed accreting systems, suggests that our understanding is far from complete. The generally accepted r-mode model would lead to a large number of observed systems in fact being unstable. This is an obvious problem that needs to be addressed by improving our models. It is, however, not clear what the missing piece of the puzzle may be.

Possibly in contrast with the f-mode, the r-mode is expected to saturate at low amplitudes due to nonlinear mode-coupling. The upshot of this is that the associated GWs are unlikely to be observed from outside our galaxy [222]. The results also imply that the spin evolution of a NS with an unstable r-mode (at saturation) may be rather complex, making a GW search even more challenging.

2. Asteroseismology

In order to understand the observed NS phenomenology we need to account for much extreme physics, many aspects of which are poorly constrained. In fact, many relevant issues will never be tested in terrestrial laboratories. Consider, for example, the possibility of quark deconfinement at high densities. While colliders like the LHC at CERN and RHIC at Brookhaven probe the properties of quark-gluon plasma, they can not reproduce the high-density/low temperature environment of a NS core. By exploring NS physics (making “sense” of observations) we can hope to constrain theoretical physics in many useful ways. In fact, this is an exciting promise of GW astronomy. The basic idea is simple. If we observe GWs from an oscillating NS, then we can use the data to infer the state of matter in the star’s core. This prospect is particularly exciting as it provides a probe of the high-density region, not the surface (where most electromagnetic phenomena arise). Of course, there is a downside to this as well. It means that we need to con-

struct models that faithfully represent the core physics. This is far from easy.

Due to their complex interior structure, NSs have many (more or less) distinct families of oscillation modes. Roughly speaking, one can associate different mode families with different pieces of physics [267]. Pressure gradients lead to the acoustic p-modes, composition (or thermal) stratification leads to g-modes, rotation leads to inertial modes, the dynamic spacetime leads to w-modes. There are modes associated with the crust, superfluidity, the magnetic field and so on. Of course, this means that the spectrum of a real NS is tremendously complicated and it may be very difficult to make sense of any data. However, for GW astronomy the situation may not be too bad, because most of the possible modes are unlikely to be efficient emitters of gravitational radiation. There are, essentially, two questions. Are there viable astrophysical scenarios where the oscillations of a star are excited to a large enough amplitude that the associated GWs may be detected? If so, what can we learn from such observations? So far, most research in this area has focussed on the second question. It has been demonstrated that global properties, such as mass and radius, can be constrained by observing f-modes (perhaps in some combination with p-modes and w-modes) [268–270]. It has also been shown that the rotational deformation, which may have a severe effect on the mode spectrum, can be “filtered out” [271]. Thermal g-modes have been studied for proto-NSs [220], and superfluid modes have also been considered [272], but we do not yet have sufficiently realistic models that we can consider the combination of the different effects. However the relevant theory framework has been developed, so it is just a matter of time until the models we consider can be considered (at least moderately) realistic.

As far as realistic astrophysical scenarios are concerned, we know that isolated NSs suffer violent events like pulsar glitches or magnetar flares. The energetics of these events is such that they could plausibly be relevant for GW astronomy. The key question is whether sufficient energy is released gravitationally. To establish this, we need to develop models that account for the observed phenomenology. Following the exciting discovery of quasiperiodic oscillations (QPOs) in the tails of giant magnetar flares [273–276], likely heralding the era of actual NS seismology, there has been significant activity aimed at understanding the dynamics of these events. Magnetars are strongly magnetized, slowly spinning NSs that exhibit high energy emission, occasionally punctuated by bursts. The favoured model contends that the energy of the magnetic field powers the observed activity, and is responsible for the bursts and occasional giant flares in soft-gamma repeaters and anomalous X-ray pulsars [277]. The observed QPOs have a complicated oscillation spectrum [278–282], the analysis of which may constrain both crust physics and the magnetic field structure [283–286]. However, the fact that these features are coupled makes the problem non-trivial. Perhaps opti-

mistically, one may assume that GWs are also generated by the large-scale, dynamical rearrangement of the core magnetic field [287–290]. However, recent numerical simulations of magneto-hydrodynamic instabilities [291] suggest that there would not be any observable gravitational radiation [292] unless the magnetic field is unphysically large [293]. In fact, the detection of GWs from magnetars in the near future seems unlikely when based on triggering f-modes in the NS [290, 292].

The recurrent pulsar glitches are also interesting, especially since they set a relatively low energy threshold for events that happen regularly in our galaxy. The general picture is that smaller glitches may be due to crust cracking, while the largest observed events are due to a transfer of angular momentum from a superfluid component to the crust (to which the magnetic field is anchored). These events could plausibly generate GWs as well, although to make definite statements about such signals is very difficult. This is not surprising, since the underlying glitch mechanism is not well understood. Available estimates range from pessimistic, suggesting that the radiated GWs will never be detected [294], to (overly) optimistic, where the signal would be borderline detectable with the first generation detectors [295]. It is quite easy to point to the flaws of each model, but to fix the relevant issues is not so straightforward. To make progress we need to improve our understanding of superfluid dynamics.

A closely related problem concerns the tidal interaction in a binary inspiral. It has been argued that the deviation from point-mass dynamics may be detectable at the late stages of inspiral [283, 296, 297]. Tidal stresses may also crack the crust, possibly leading to an electromagnetic signal that would precede the merger [298, 299]. This problem is interesting, and it requires the same computational technology as the seismology problem. Again, the challenge is to build truly realistic NS models, and assess the impact of the many different pieces of physics involved in the dynamics of the star. Possible electromagnetic precursors are also of obvious interest for GW searches.

III. DETECTORS AND THEIR CAPABILITIES

The observation of GWs, combined with astronomical observations from gamma-ray and X-ray satellites, optical/radio telescopes, and neutrino detectors, will enable a new, comprehensive multi-messenger astrophysics which will play a transformative role in our understanding of the Universe, with new constraints to source models combined with identification of the host galaxy, redshift and luminosity distance. In this section, we review present and prospected instrumental capabilities for multi-messenger observations in the foreseeable future.

A. Gravitational-Wave Interferometers

The inception of ground-based interferometers represented significant progress toward the detection of GWs. In 2005-2010 the Laser Interferometer Gravitational-wave Observatory (LIGO) [14] and Virgo [15] operated four detectors, sensitive to the merger of two NSs within ~ 30 Mpc of Earth [300, 301]. A GW interferometer uses lasers to monitor the relative distances between the beam splitter and mirrors located at the end of its two arms. A GW signal will stretch one arm of the interferometer and compress the other, causing a detectable change in the interference pattern at the output of the interferometer. First-generation GW interferometers were capable of observing a change in the arm length of 10^{-18} m, or about 1/1000 the diameter of a proton, around the most sensitive frequency (~ 100 Hz). Despite this impressive sensitivity no GWs have been detected so far, due to the low expected event rates, as discussed in Section II A 3 for the merger of compact binary systems.

The second generation of GW interferometers, scheduled to start within this decade, will have improved seismic isolation, suspension, optics and laser systems, offering roughly a factor 10 sensitivity improvement over a wide frequency range [2]. These detectors could allow us to search for GWs from the rich class of transients discussed in Section II.

In general, transient searches for GW signals fall into two categories: modeled and unmodeled (or weakly modeled). Modeled searches rely on the availability of precise template waveforms for some signal classes, such as the known inspiral waveforms from compact binaries, against which data can be compared with matched filtering techniques [302]. Unmodeled searches can use the coherence between excess power in multiple detectors to distinguish signals from noise [303]. Therefore, it is critical to have several detectors with comparable sensitivities operating with a high-coincidence duty cycle.

A network of detectors brings significant resistance against nonstationary noise due to environmental and instrumental disturbances, as well as sensitivity in a greater volume [303, 304]. Two advanced LIGO detectors [2] in Livingston, LA and Hanford, WA, and one advanced Virgo detector [3, 15] are currently under construction and will start operation in a few years. A large cryogenic GW telescope, KAGRA, [9, 305] is being constructed in Japan, and a third advanced LIGO detector is currently under consideration for being built in India [8]. The LIGO and Virgo collaboration have released a plan for the commissioning deployment of a network of second-generation GW detectors, which will start in 2015 with short (few months) science runs with two detectors, and will grow to stable operation of a network of 4 detectors by 2022 [306].

The GW detector network performance greatly depends on the number of detectors in the network, their geographical location, and the relative orientation of the detector arms. For each direction in the sky, the perfor-

mance of the detectors is characterized by their antenna patterns, which can be combined into the network antenna factor [303]. For example, Figure 2 (left plots) shows the distribution of the network antenna factor as a function of the sky coordinates. Note that the six-detector network provides more uniform coverage of the sky than the Hanford-Livingston or Hanford-Livingston-Virgo networks. The right-hand panel of Fig. 2 shows the alignment factor between detectors in the network; networks of few detectors with similar arm orientations may only be sensitive to one of the two GW polarizations for some sources.

A worldwide GW detector network enables the reconstruction of source parameters and the localization of GW events on the sky [303, 307–311]. Accurate source localization is key to enabling multi-messenger astronomy, e.g., via joint observations with electromagnetic telescopes [312–315]. Meanwhile, measurements of other source parameters can enable studies of astrophysics and tests of general relativity [96, 316–318]. Reconstruction of source location and polarization requires geographically-separated detectors to make independent observations of the same GW event; recovery of intrinsic parameters such as binary component masses by modeled searches primarily depend only on the total network SNR [311].

Proposed future GW instruments will further enhance our astrophysical reach. These include third-generation ground-based detectors [319–322] that would be an order of magnitude more sensitive than second-generation instruments. Meanwhile, pulsar timing arrays [323] would be sensitive to binaries composed of the most massive black holes in the Universe, while the Laser Interferometer Space Antenna or a similar space-based detector [12, 13] could detect massive black hole mergers, extreme mass ratio inspirals, and tens of thousands of Galactic white dwarf binaries.

B. Gamma-Ray and X-Ray Instruments

The continued operation of high-energy, electromagnetic satellites in the coming decades is essential for the realization of the full science potential of joint GW-GRB observations discussed in Section II. A GRB/X-ray trigger, possibly followed by optical observation to identify the host galaxy and source redshift, can be used as input for a GW transient search at a known time and location. In turn, the worldwide network of GW observatories will be able to reconstruct in near real time the sky position of GW candidates and trigger multi-messenger observations for apparent afterglows. The electromagnetic observations may identify the source’s host galaxy, redshift and, assuming standard cosmology, the luminosity distance, while the GW measurement may offer enhanced constraints on the source engine; a joint observation will ultimately enable us to decipher science that would otherwise be inaccessible.

Currently, it is hard to know which GRB and X-ray sensitive instruments and all-sky transient surveys will be operating in the next two decades. Present GRB transient observation satellites such as AGILE [324], Fermi [325], INTEGRAL [326] and Swift [327] are not guaranteed to operate in the era of Advanced LIGO and Virgo. It is possible that the lifetime of some of these instruments will be extended, unless an operational constraint or accidental equipment failure necessitates end-of-life procedures for the satellites.

Ongoing X-ray missions such as Chandra [328] and XMM-Newton [329] are expected to continue to provide information on GRB properties by observing their X-ray afterglows, at least during the early years of the advanced GW detector era, but do require prior localization of the GRB. The Suzaku [330] orbiting X-ray observatory has a capability of detecting GRBs via its wide-band all-sky monitor. The recently launched NuSTAR [331], targeting the hard X-ray region, is likely to operate much beyond its originally planned lifetime of two years. MAXI [332] is an all-sky X-Ray monitor, installed on the International Space Station by the Japan Aerospace Exploration Agency [332].

New projects include the Indian satellite ASTROSAT [333–335], a multi-wavelength mission that will monitor the X-ray sky for new transients and is currently scheduled for a 2013 launch, with an expected lifetime of 5 years. The next Japanese X-ray Astronomy mission, ASTRO-H [336], is scheduled to be launched in 2014; it can contribute to follow-up observations of GRB afterglows at high resolution [337]. The launch of SVOM [338–340] – a joint Chinese-French GRB monitor mission with an extended spectral coverage from the visible to a few MeVs, and with good GRB localization capability – is planned towards the end of the decade.

In addition, there are numerous high-energy astrophysics mission concepts targeting the X-ray and gamma-ray spectrum [341]. Some (e.g. AXTAR, EXIST, Xenia, GRIPS, A-STAR, JANUS) would allow the detection of X-ray or gamma-ray transients with potential GW burst counterparts. NASA’s X-ray Mission Concept Study Report [342] states that it is feasible to start a next X-ray mission toward the end of the decade. In the study report, simplified missions that capture most of the recently terminated IXO (International X-ray Observatory) mission science goal elements were identified. The Lobster Transient X-ray Detector is a mature concept that was proposed to be deployed on the International Space Station in three to four years. Its unique technology would allow to detect transient X-ray emissions from a large portion of the sky at a wide field of view and high sensitivity. If approved, the instrument could work in conjunction with ground-based GW detectors, following up GW candidate events [343].




FIG. 2. The distributions of the network antenna factor (left plots) and the network alignment factor (right plots) as a function of latitude (θ) and longitude (ϕ) for the Hanford-Livingston network, Hanford-Livingston-Virgo network and a six-detector network (from top to bottom).

C. Electromagnetic Instruments from Radio to UV

Low-latency electromagnetic follow-up observations of GW event candidates will enable the identification of possible optical and other electromagnetic counterparts [90, 344–348]. The infrastructure for this type of analysis was tested during the most recent LIGO-Virgo data run, in 2010, when observation requests were sent to wide-field optical telescopes and other instruments, including QUEST, TAROT, ZADKO, Pi in the sky, ROTSE, SkyMapper, LOFAR, the Palomar Transient Factory [312] and the Swift satellite [313].

Radio telescope arrays, such as LOFAR [349], EVLA [350], ASKAP [351] and the future Square Kilometre Array [352], have in most cases a wide field of view and are able to provide sub-arc-second angular resolution, that is superior to the pointing of the advanced GW detector networks of the future.

In some theoretical models various mechanisms may give rise to a prompt pulse (see e.g. [90]), strong winds or bursts [61], flares [298, 299], intense Poynting fluxes

and emissions through shocks [62, 353] or afterglow radio emission (see e.g. [354]) from some expected GW sources, particularly coalescing compact binaries, thus motivating coordinated observations. In addition, the use of GW detectors as a trigger for follow-up radio searches could provide a method of detecting faint radio transients that might otherwise be missed.

Prospects of electromagnetic counterpart observations of GW events were recently discussed in [355–360]. Gamma-ray observations will be critical for confirming a connection between short GRBs and NS-NS/NS-BH mergers. Optical and radio afterglows, even off-axis, as well as r-process nucleosynthesis “kilonovae” are detectable in principle, provided that an optimized search strategy is used, taking into account the emission timescale and instrument parameters.

D. Low-Energy Neutrino Detectors

Core collapse supernovae were discussed in Section II C as sources of GW radiation and engines for long gamma-ray bursts. However most of their energy, about 99% of $\sim 10^{53}$ ergs, is released in the form of neutrinos, within a few tens of seconds immediately following the collapse. The neutrinos in this burst are of all flavors, and their energy ranges from a few to tens of MeV. This prediction was confirmed by the detection of a burst of 19 neutrinos from SN1987A in the Large Magellanic Cloud by two water Cherenkov detectors: IMB in the United States [361] and Kamiokande II in Japan [362]. Scintillation detectors also reported observations [363, 364], and the main features of the signal [365, 366] confirmed the baseline model of stellar collapse (see e.g. [151, 187]).

Several existing neutrino detectors are sensitive to a neutrino burst from a galactic supernova [367, 368]. Super-Kamiokande [369], a 50-kiloton water Cherenkov detector in Japan, would observe ~ 8000 events from a supernova ~ 8.5 kpc away [370]. The LVD [371, 372] and Borexino [373, 374] scintillation detectors at Gran Sasso in Italy, KamLAND [375, 376] in Japan, and the upcoming SNO+ [377] in Canada would also observe hundreds of neutrino events interacting in 300-1000 tons of liquid scintillator. The IceCube detector [378] is a cubic-kilometer detector located at the geographic South Pole. IceCube is nominally a multi-GeV neutrino detector, but it is also sensitive to MeV-neutrinos from a Galactic supernova and could observe an increase in the count rate due to a diffuse burst of Cherenkov photons in the ice [379]. Super-Kamiokande, LVD, IceCube and Borexino operate as part of the SNEWS (SuperNova Early Warning System) network [380, 381], for a prompt alert to astronomers in the case of a supernova neutrino burst.

The distance reach of the global network of neutrino detectors covers the Milky Way and a significant fraction of its Satellite System, up to $\sim O(100)$ kpc. This can be considered as a good match to the GW network reach for core-collapse supernovae [382–386] of GW detectors. A 50-kiloton detector like Super-Kamiokande has a very low chance of detecting a SN in M31 (at ~ 770 kpc) on its own. Low-energy neutrino detectors have angular resolution comparable to advanced GW detector networks [307], and this makes synergetic observations particularly desirable [387, 388].

Long-term plans call for neutrino detectors with up to ~ 5 megatons of fiducial mass [389, 390] to enable observation of neutrinos from M31 and M33 [391, 392]. Proposals include MEMPHYS [393], LENA [394], and GLACIER [395] in Europe, DUSEL LBNE [396] in the US, Hyper-Kamiokande [397] and Deep-TITAND [391, 398] in Japan. It is not unreasonable to expect that the lifetime of these detectors will coincide with the operation of third-generation GW detectors.

E. High-Energy Neutrino Telescopes

High-energy neutrinos (HENs) in the GeV–PeV range could also unveil new physics in joint observations with electromagnetic and GW signatures [399, 400]. The detection of HENs is pursued in large Cherenkov detectors that exploit their charged-current interaction in large volumes of water or ice. Most of the neutrino energy is transferred to a single high-energy electron, muon, or tau particle, which will emit Cherenkov radiation as it travels through the detector medium. High-energy muons are most useful for neutrino astronomy, since they don’t lose energy as rapidly as electrons and have longer lifetime than taus, so their path can be several kilometers long. The Cherenkov light emitted along this path can be detected and used to measure the direction and energy of the muon, and thus of the primary neutrino.

There are three HEN observatories currently in operation. The IceCube observatory [378] has recently been extended with an additional component called DeepCore [401], designed to be sensitive to neutrino energies below IceCube’s lower limit of ~ 100 GeV, down to 10 GeV, effectively increasing the detector’s astrophysics reach [402]. ANTARES [403], located in the Mediterranean sea, is scheduled for an upgrade to a cubic-kilometer detector called KM3NeT in the coming years [404]. A third HEN detector operating at lake Baikal is also planned to be upgraded to a km^3 volume [405].

The distance reach of high-energy neutrino detectors is virtually infinite, although at larger distances the probability of detecting a neutrino from an individual source diminishes. The Waxman-Bahcall model [406], the benchmark model of HEN emission from GRBs, predicts about $n_{\text{HEN}} \approx 100$ neutrinos detected in a km^3 detector for a typical GRB at 10 Mpc [407], although recent upper limits from the IceCube detector disfavor GRB fireball models with strong HEN emission associated with cosmic ray acceleration [408]. However, milder HEN fluxes or alternative acceleration scenarios are not ruled out [409]. Moreover, the constraints weaken substantially when uncertainties in GRB astrophysics and inaccuracies in older calculations are taken into account, and the standard fireball picture remains viable [410, 411].

Models of high-energy neutrino emission from mildly relativistic jets of core-collapse supernovae, and potentially from choked GRBs, predict HEN emission of $n_{\text{HEN}} \approx 10$ [412] (note that the result presented in [412] is three times higher, as it does not take into account neutrino flavor mixing). Horiuchi and Ando [413] estimate n_{HEN} from reverse shocks in mildly relativistic jets to be $n_{\text{HEN}} \approx 0.7 - 7$ for a km^3 neutrino detector (after taking into account neutrino flavor mixing). Razzaque *et al.* [414] obtain $n_{\text{HEN}} \approx 0.15$ for supernovae with mildly relativistic jets with jet energy of $E \sim 10^{51.5}$ erg.

The most stringent observational constraints on transient GW+HEN sources so far has been obtained using searches with the latest Initial LIGO-Virgo (S6-

VSR2/VSR3) detectors and the IceCube detector in its 40-string configuration [415]. The derived constraints were also used to estimate the science reach of the Advanced LIGO-Virgo detectors in combination with the completed IceCube detector, with promising results. The first joint search of ANTARES, LIGO, and Virgo data for coincident GW and HEN using LIGO, Virgo and ANTARES data derived limits on the rate density of joint GW-HEN emitting systems in the local universe, comparing them with densities of merger and core-collapse events [416] that are compatible with previous results. While the available upper limits for initial detectors impose no constraints on joint emission models, the results [415] show that advanced detectors will be able to constrain some emission and population models, therefore also having the potential of detecting joint sources.

IV. CHALLENGES AND OPEN QUESTIONS

The initial generation of GW detectors has targeted most of the transient sources described in Section II, yielding observational limits which already bear astrophysical interest. For instance, for two nearby GRBs, the non-detection of a GW signal made it possible to exclude a merger as the GRB central engine [417, 418], while constraints on GW emissions were produced for magnetars [419] and in a study of the 2006 Vela pulsar glitch [420]. Population constraints have been produced for GWs in coincidence with gamma-ray bursts [421], and all-sky limits have been set on rates of binary mergers [301, 422, 423] and generic bursts [424]. A first coincidence search with high-energy neutrinos has been performed [416], and in the most recent data run (2009-2010) transient candidates have been broadcast for EM follow-up [312, 313]. The potential for extracting fundamental physics and astrophysics from GW data should be dramatically enhanced by the next generation of GW detectors, with a projected ~ 1000 times larger sensitive volume and the newly accessible $\sim 10 - 40$ Hz frequency band. Predictions for compact binary detection rates with next generation detectors, based on astrophysical observations, population synthesis and source models, are available in [101].

The engagement between the experimental and theoretical communities in the coming years will shape the future of GW astrophysics: to maximize their scientific output, GW transient searches will need to include more information from theoretical and computational astrophysics. In turn, robust source modeling will be required to provide a theoretical understanding of the mapping between signal characteristics and physics parameters, including the knowledge of potential degeneracies that may be broken by complementary information from multi-messenger observations. In this section we elaborate on what we identify as the main open challenges in the theoretical understanding of GW transient sources and in the ability to identify their signature.

A. Observations of Gravitational-Wave Transients

Despite steady progress in relativistic astrophysics, there is still significant uncertainty in predicted waveforms for most GW transient sources. In some special cases, when an accurate signal model is available, the search relies on matched filtering with a bank of templates [301, 422, 425, 426]. In the more general case, GW bursts can be identified in the detector output data as unmodeled excess power localized in the time-frequency domain [424].

The principal challenge for the instruments is to achieve a 10-fold improvement in sensitivity compared to the first generation of GW interferometers, including a reliable calibration and a low rate of noise transients [e.g., 427, 428]. As soon as the second generation of GW detectors reaches design sensitivity, the main experimental challenge to GW burst science will be to discriminate between real GW signals and noise transients that happen to coincide in multiple detectors.

A fourth site, in addition to the existing facilities in the USA (LIGO Hanford and LIGO Livingston) and Italy (Virgo), is an additional instrumental challenge that will enable source localization and increase detection confidence. As discussed in Section III, the identification of a consistent signal in a network of instruments is a key ingredient in GW searches, dramatically increasing the confidence in a candidate event. This is especially important for a GW burst, which otherwise may not be distinguishable from noise fluctuations of instrumental or environmental origin. A statistic built from the coherent sum over the detector responses is used to rank candidate events and discriminate between signal and noise, yielding better sensitivity (at the same false alarm rate) than individual detector statistics. Accurate timing information from multiple detectors also makes it possible to reconstruct the two GW polarizations and localize the source on the sky. The LIGO and Virgo collaborations outlined a timeline for the localization of GW transients by advanced detectors [306], in order to facilitate the formulation of joint detection strategies [429] with electromagnetic, neutrino, or other observing facilities.

The level of background, and, hence, the significance of a candidate event, is estimated empirically with time slides. Tests with simulated signals injected into LIGO and Virgo noise show that when a signal model can be assumed, this technique makes it possible to achieve high detection confidence. For instance, in a detection exercise during the most recent LIGO-Virgo run, the simulated coalescence of two compact objects with network signal-to-noise ratio of 12.5 was identified with a false alarm rate of 1 in 7000 years using matched filtering [301]. A completely un-modeled search instead yielded $1/1.1$ years, after accounting for trial factors [424].

An important challenge for GW burst searches in the advanced detector era will be to incorporate partial information from theoretical models in order to constrain the false alarm rate and increase detection confidence in

analyses tuned to specific sources, with perfectly modeled and completely general searches as extreme cases. Understanding what information can be incorporated in searches, and learning how to make inferences on source parameters from transients, starting from the frequency, bandwidth and duration of a coherent event, will require a close synergy with the modeling community. One particularly exciting possibility is the use of GW signals, possibly in coincidence with electromagnetic observations, as probes of strong-field dynamics and tests of general relativity itself [e.g., 430].

We highlighted three key areas of cooperation for the relativistic astrophysics community:

1. *What can we understand about the source from the data?* Theorists and modelers can guide targeted analyses for a specific source, interpret the astrophysics of general searches, and (together with instrumentalists) determine what science can be extracted from the detector response. In return, the data analysis community should instruct astrophysicists/modelers on how to combine detector capabilities and theoretical waveforms into a prediction about the physics that can be extracted from a detection, and a realistic assessment of the extent to which these predictions are reliable.

2. *Can we improve the pipeline that moves theoretical advances in source modeling into the GW data analysis search methodologies?* Both theoretical and computational understanding of GW sources and GW search methodologies are rapidly advancing. There is, however, a time lag in implementing some of these advances into actual LIGO searches. One of the challenges in anticipation of the advent of GW astronomy is the development of better strategies to rapidly get the best models to data analysts.

3. *How to bring together GW, EM and astroparticle physics?* Searches for GW bursts in the advanced detector era will be a combination of untriggered, all-sky searches and externally triggered, localized searches. All-sky searches have the greatest potential for finding electromagnetically dark sources, and may discover unexpected signatures. They also provide triggers for follow-up studies of candidate events with EM observations. Externally triggered searches will have EM and/or neutrino counterpart observations. Strategies are needed to combine EM, neutrino, and GW information and foster collaborations between the GW community and “traditional” astronomers, from data sharing to physics extraction.

B. Double Neutron Star and Neutron Star – Black Hole Binaries and Short Gamma-ray Bursts

While the most widely accepted model for short GRB progenitors is the merger of a NS with another NS or a BH [27], Swift and Fermi have observed only a few short GRBs per year, and therefore the prospect of observing a

short GRB in association with a GW counterpart appears challenging. Furthermore, it is not easy to identify the host galaxy of a short GRB because the sky localization is poor unless an afterglow is detected, and even then, natal kicks could displace the binary system from the host galaxy [22, 431]. The nearest short GRBs may be the hardest to associate with their host, since angular offsets could be large. This uncertainty in host galaxy identification leads to an uncertainty in the progenitor distance, which, in turn, leads to an uncertainty in GW signal association. However, given the relatively small reach of advanced GW detectors, it is possible for GW observations to rule out potential host galaxies that are too distant.

There are many outstanding questions about the nature of observed GRBs. It is unclear how to best distinguish short GRBs associated with merging compact binary progenitors from soft gamma-ray repeater hyperflares. The energy budget for short GRBs is also unclear, although they have been identified (via their host galaxies) up to high redshifts. In particular, there are several uncertainties about the beaming of short GRBs, with very few measurements up to now [26]. Beaming is also a crucial ingredient to understand how many GW events could be associated to the observed rates of short GRBs (for a recent review of the properties of short GRBs, and the implications for their progenitors, see [432]).

There are several open problems in the astrophysics of the merging systems that are very important in the modeling of compact binary mergers. If magnetic fields are indeed responsible for the energy budget, where do these large fields come from in a merger scenario? The magnetic fields of the progenitor NSs are not large enough and the merger event can amplify them. Is this amplification sufficiently strong? Does the spin-down of the final BH cause any late emission signature? Shall we be able via more complete modeling to use GW or EM observations to distinguish between a NS or a BH as the companion compact object in the merger?

Observing GWs from compact binaries also challenges our understanding of binary evolution. If we can better understand merger events, we may be able to use GW and EM observations to understand physical mechanisms that are responsible for compact binary formation and the corresponding distribution of observed events. Another challenge is posed by the poorly known nuclear physics at the densities typical of NS interiors. Do we understand the physics of nuclear processes at NS densities well enough to accurately predict GWs from these systems? Even with an improved understanding of the properties of matter within NSs, adding an accurate treatment of magnetic fields, together with realistic NS EOS and neutrino radiation to get the right post-merger dynamics will be a significant technical and computational challenge to the numerical relativity community. It will be especially important to be able to efficiently generate a large number of data sets to be compared with GW observations. While the GW inspiral signal could prob-

ably be modeled analytically (e.g. using variants of the effective one-body model [53, 55]) facilitating the production of a large number of templates, the merger and post-merger dynamics will require a large and accurate set of numerical simulations.

C. Binary Black-Hole Mergers as Transient Sources

Numerical relativity provides direct solutions of the Einstein equations for binary BH spacetimes. Numerical relativity plays a major role in two large collaborations using binary BH waveforms. The NRAR (Numerical Relativity/Analytical Relativity) collaboration is working to provide templates that cover the inspiral, merger and ringdown phases of binary BH coalescence. The NINJA (Numerical INjection Analysis) project [433, 434] is using hybrid numerical relativity waveforms to test data analysis algorithms for the detection and characterization of GWs. The hybrid waveforms are made by stitching post-Newtonian and numerical relativity waveforms together. While the waveforms and templates created in these projects are useful for burst analysis, and in the case of NINJA unmodeled burst search algorithms are being tested, a burst-only focused study offers an opportunity and challenge to theory and data analysis.

For stellar-mass binary BH systems the challenge is to provide template coverage for modeled searches over a large parameter space using a combination of post-Newtonian techniques and numerical relativity. This includes the challenge of stitching the post-Newtonian and numerical waveforms together. The choice of where to make the stitching (i.e., how many cycles are supplied by post-Newtonian theory and how many cycles must be simulated numerically) depends on the physical parameters of the binary. The NRAR and NINJA collaborations are working systematically to address these problems.

More massive systems offer a different challenge to the numerical relativity and data analysis communities. Unmodeled or weakly modeled burst searches are most appropriate when we know little about the signal model. Rather than the exquisite accuracy in phase necessary for template building, a burst-only focus in numerical relativity offers the opportunity to survey the physical parameter space of binary BHs in a timely manner by reducing the resolution required compared with inspiral-focused simulations. The challenge, however, is learning to identify key features of the merger that will be useful to the data-analysis algorithms for transient sources, potentially improving glitch identification and reducing false-alarm rates. Only one of these studies, aimed at investigating the sensitivity of the Omega burst algorithm [435] to changes in the BH spin vector [436], has been completed, and more studies of this kind are needed.

Prior to the first science runs of the advanced detectors, we need information about the merging systems, including the energy, angular and linear momentum ra-

diated (in total and in each harmonic mode), the polarization (such as ratios of circular to linear) and mode dependent merger time-scales. This provides great opportunities: can we increase the likelihood of detecting GWs from binary BH mergers by employing weakly modeled burst searches? Can information gleaned from NR waveforms of merging BHs reduce the false-alarm rate for burst searches? Can a burst algorithm find sources of merging BHs for physical parameters that have not yet been modeled, or have not been modeled well enough? In order to answer these questions, we need to determine the smallest number of cycles we can simulate in numerical relativity while still representing actual signals reasonably well. We need to figure out how the inevitable lack of phase accuracy or length of numerical simulations propagates in the search for a transient source. This leads us directly to perhaps the most interesting and important challenge for a transient search of the binary BH merger sky: how well can such a search determine the physical parameters of the merging sources? Perhaps a burst-first approach (targeted at first finding a burst signal, and then digging through the data with matched-filtering parameter estimation techniques) can help in this regard.

D. Core-Collapse Supernovae and Long Gamma-ray Bursts

Uncertainties abound in current GW estimates from core-collapse supernovae. These uncertainties include poorly known initial conditions, the relevance of 3D effects in the explosion mechanism, or the impact of detailed 3D neutrino transport on the GW signal from anisotropic neutrino emission. Here we focus on two major issues in determining accurate GW signals: rotation of the progenitor and convective instabilities.

1. The Role of Rotation.

Rapidly rotating systems have the potential to produce strong GW signals, especially if nonaxisymmetric instabilities develop. Whether or not such signals occur in nature depends upon the rotation profile of the pre-collapse star. Classic models of single stars produce a range of core rotation speeds, depending upon the initial angular momentum of the star and the magnetic field generation [166, 437–441]. In the absence of magnetic fields, the core can decouple from the rest of the star, retaining a high spin rate even as the stellar envelope expands and decreases its spin rate. But if magnetic fields are produced at boundary layers, they tie the core to the stellar envelope, causing its spin to decelerate with the envelope.

The high-spin requirements of most long GRB progenitors have pushed stellar modelers to seek new ways to produce fast-spinning cores. Fast-rotating single star models often invoke extended mixing and generally re-

quire low metallicity, below $\sim 0.1Z_{\odot}$ [440–442]. Others have devised a number of binary progenitors of long GRBs that may also produce fast rotating cores invoking tidal interactions, mass transfer, and common envelope mergers [443–447]. The fastest-rotating cores collapsing to form NSs could be produced in the merger of two white dwarfs [448–453], but we currently do not know whether such systems produce thermonuclear or core-collapse supernovae.

Uncertainties in the evolution of massive stars (especially rotating stars) and in binary evolution make it very difficult to predict the distribution of spin rates of collapsing cores. If fast-spinning pulsars also develop strong magnetic fields, selection biases make it difficult to use the observed pulsar distribution to constrain the birth spin rate of NSs. Similarly, high spin rates are just one requirement for long GRB engines, and extrapolations to the spin rates of newly formed NSs from long GRB rates can only be done with specific model assumptions.

2. Convective Instabilities.

The detailed nature of the convective instabilities (both within and above the PNS) also remains an open question in estimates of GW emission. If low-mode large-scale convection develops and is modulated by the SASI, as is borne out by current simulations, we have a fairly strong picture of the GW signal. But the nature of this convection (and of the resulting GW signal) is different between 2D and 3D models. The current resolution in core-collapse supernova simulations, in particular in the first 3D simulations, remains roughly an order of magnitude lower (per dimension) than what computational fluid dynamicists argue is required to accurately model convective modes and turbulence [454, 455]. In addition, with few total turnover timescales in the model, initial perturbations (caused by convection in the Silicon burning layer) may have a strong impact on the convective motions. To more reliably predict the GW signal from convection, it will be necessary to gain a better understanding of convection, the SASI, and of the impact of perturbations from late convective burning stages.

E. Isolated Neutron Stars

It should be clear from our discussion that transients in isolated NSs provide a range of interesting prospects for GW astronomy. At the same time, it must be appreciated that the expected signals are likely to be weak. Because of the complexity of the physics involved in most realistic scenarios, these signals are also extremely difficult to model in detail. This means that it may be unrealistic to expect modeling to produce accurate search templates for matched filtering. Dealing with this difficulty is one of the main challenges for the next few years. Realistically, the sources that we have discussed here may

only be borderline detectable with second-generation detectors. They may require future developments, like the Einstein Telescope, with enhanced sensitivity in the high-frequency regime. However, such detectors could provide us with an unprecedented view into NS interiors [268, 269, 271, 456], especially if the signals are strong enough that the asteroseismology strategy is viable. The hope is that GW observations will complement EM data, helping us to place constraints on physics under extreme conditions.

On the theory side, it is easy to identify directions in which the current models need to be improved. Basically, we need to continue developing both the computational technology and our understanding/implementation of the relevant microphysics. In order to model the relevant astrophysical scenarios, we obviously need to build more realistic models. This is associated with a number of challenging issues. For the purposes of this summary, let us consider two specific problems.

First of all, it is clear that magnetars may be interesting target sources given their, sometimes rather violent, activity. However, even though there has been clear progress in recent years, we are still quite far from modeling “realistic” magnetized stars. The difficulties range from fundamental to technical. We do not yet have reliable models of compact magnetic equilibria with the expected composition, and we must account for superconductivity (and the associated magnetic flux tubes) in the star’s core.

Secondly, recent work on instabilities, such as those of the r-mode, has shown that we do not, in fact, understand the relevant dissipation mechanisms well enough to explain the observed population of fast spinning, accreting NSs. Given the amount of effort that has gone into modeling the r-mode instability, this is somewhat embarrassing. Another major problem concerns the nonlinear evolution of this kind of instability. The associated timescale is too long for the problem to be within reach of nonlinear simulations. At the same time, we need to account for nonlinear aspects (like the mode-coupling that is thought to saturate the r-mode). This is truly problematic, especially if we want to make the stellar models more realistic by including the appropriate microphysics. Similar issues relate to pulsar glitches, with a relaxation time of weeks to months but an initial rise time of tens of seconds, and NS mountains, which may evolve due to plastic flow.

These problems are clearly far from trivial, but they provide a good illustration of the kind of issues that we need to deal with if we want to extract as much physics as possible from future GW signals from NSs.

F. Multi-Messenger Astrophysics of Transient Sources

The success of the GW burst endeavor relies on a continued dialogue between source theory and detector ex-

pertise, or between predicted sources and what can be achieved in a practical instrument. On one side this drives detector design, on the other it determines which parts of parameter space theorists should focus on.

While GW bursts generally do not have well-defined waveforms, they have robust signal features when characterized by their energy. The duration, central frequency or even time-frequency characteristics of the signal energy can be used to tune source-specific searches. A GW burst search can be informed by source population or emission models to identify relevant regions of the parameter space to focus over. Moreover, any waveform estimates from potential sources can be used to characterize the efficiency of the developed searches. When a population of sources or repeating GW emitters are considered, it is possible to design an analysis to target the energy features to build up SNR and reject noise events.

While all-sky, untriggered GW searches are the principal mode of operation for GW detectors, triggered searches based on observations in other bands or theoretical guidance on the most important potential sources can yield to improved sensitivity. We will want to revisit the issue of optimal sources for triggered GW searches as new astrophysical discoveries are made and GW detector performance and data analysis capabilities improve [e.g., 359]. Meanwhile, once potential GW sources have been identified, to facilitate efficient observations, it is highly desirable to exchange triggers between the GW and astronomy communities within 1 hour of the trigger, or even less.

Research on GRBs is now a mature subject. There are redshifts for hundreds of GRBs (all extragalactic). Despite this, many uncertainties remain. What is the nature of their progenitors? At least some long GRBs are confidently associated with CCSNe, but what about the other observations? GRB emission processes (both non-thermal and thermal) are quite uncertain, and the total energy budget is unclear.

A lot of infrastructure has been put in place and communication channels between the GW and astronomy communities opened in the preparation for the Advanced LIGO era. It is worth asking, in the multi-messenger era, how should this momentum be maintained? A post-Swift mission with a comparable source localization error box will complement GW observations. Facilities with wide-field search capability (snapshot vs. tiling) and a field of view covering Advanced LIGO error regions may allow a rapid discovery of the EM transient counterpart, but it is equally important to have high-resolution instruments that can follow up the transient discovery with other observations, such as obtaining a source spectrum. There is also tension on the GRB requirements between the GW community and the broader GRB community: missions that suit multi-messenger astronomy would focus on rapid observations of closer GRBs, which may not be what other GRB astronomers (who may want to explore the oldest GRBs and cosmology) are after.

V. CONCLUSIONS AND PERORATION

The era of GW astronomy is upon us. Advanced ground-based detectors are expected to reach their design sensitivity near 2019. The space-based GW detector eLISA may be selected in the EU as one of the next L2/L3 missions, with a launch possibly scheduled before the end of the next decade. IPTA may also detect GWs in the next few years [457, 458]. This paper has focused on transient GW sources for second-generation Earth-based detectors, particularly those amenable to unmodeled or weakly-modeled burst searches.

We have reviewed the state of the art of theoretical and observational studies of several GW transient sources. Compact binaries composed of NSs and/or BHs are among the most promising sources, because they are detectable by advanced ground-based interferometers at distances of hundred of Mpc. In recent years, numerical simulations of these objects have increased our knowledge of their GW signals and, more recently, of their possible connection to EM observations. In particular, NS-NS and NS-BH binaries are thought to be linked to SGRB engines. A coincident detection of GWs and SGRBs would be the smoking gun that SGRBs are indeed caused by compact object mergers. Another GW source that has a very important connection with EM observations are CCSNe, which are currently the main candidate LGRB engines. Unfortunately, GWs generated by such systems will be detectable only for events localized in our Galaxy, hence the expected event rates are quite low.

Due to the possibility that the first GW detection may happen in the next few years, it is important to increase the current efforts in modeling GW sources. Significant progress has been made in modeling binary BHs, but covering the template space of generic precessing binaries is still a big challenge. New theoretical and computation insights are needed to tackle BHs spinning close to the Kerr limit, even for nonprecessing binaries with aligned spins. The study of non-vacuum sources, due to the higher level of complexity, still requires significant effort in order to better understand the effects of several different physical mechanisms on their evolution (e.g. EOS, neutrino emission, magnetic fields, convection, turbulence). While part of this work will require the development of better and more efficient numerical codes, several studies will also require the use of larger computational resources. For example, the study of convective instabilities in CCSNe and of magnetic instabilities in NS mergers will require much higher resolutions, and hence more powerful computational resources than those used so far. Moreover, since the properties of matter in both NS mergers and CCSNe are not yet well known, it will be imperative to build a large and publicly available database containing GW signals generated by different models. This will be crucial in order to be able to extract physical information from the GWs that will be detected from these systems.

Detecting EM counterparts to GW events will lend greater confidence to GW searches and provide useful complementary information. This will considerably help in source localization and, in the case for example of SGRBs and LGRBs, allow us to better understand the engine of some of the most fascinating astrophysical phenomena. It will be particularly important that both X-ray and γ -ray satellites will be available and ready when the second generation of GW detectors will start to take data. For these studies to be successful, it will be necessary to increase the current level of collaboration between the GW and EM communities, and ideally better coordination between funding bodies.

After almost 100 years since the prediction of their existence, the direct detection of GWs is on the horizon. In the next decades GW astronomy may become one of the most important sources of information on high-energy astrophysics, providing us with an unprecedented level of information on NSs and BHs.

ACKNOWLEDGMENTS

The “Gravitational Wave Bursts” workshops in Chichen-Itza, Mexico (December 9-11, 2009) and Tobermory, Scotland (May 29-31, 2012) were supported by National Science Foundation Grants Number 0946361 and 1231548. The authors gratefully acknowledge the support of the United States National Science Foundation for the construction and operation of the LIGO Laboratory and the Science and Technology Facilities Council of the United Kingdom, the Max-Planck-Society, and the State of Niedersachsen/Germany for support of the construction and operation of the GEO600 detector. The authors also gratefully acknowledge the support of the research by these agencies and by the Australian Research Council, the International Science Linkages program of the Commonwealth of Australia, the Council of Scientific and Industrial Research of India, the Istituto Nazionale di Fisica Nucleare of Italy, the Spanish Ministerio de Educación y Ciencia, the Conselleria d’Economia, Hisenda i Innovació of the Govern de

les Illes Balears, the Royal Society, the Scottish Funding Council, the Scottish Universities Physics Alliance, The National Aeronautics and Space Administration, the Carnegie Trust, the Leverhulme Trust, the David and Lucile Packard Foundation, the Research Corporation, and the Alfred P. Sloan Foundation. We thank the Kavli Institute for Theoretical Physics at UC-Santa Barbara, supported in part by NSF grant PHY11-25915, for hosting the workshop “Chirps, Mergers and Explosions: The Final Moments of Coalescing Compact Binaries.” The Columbia Experimental Gravity group is grateful for the generous support from Columbia University in the City of New York and from the National Science Foundation under cooperative agreement PHY-0847182. E.B. is supported by National Science Foundation through CAREER Award Number PHY-1055103. K.B. acknowledges NASA Grant Number NNX09AV06A and NSF Grant Number HRD 1242090 awarded to the Center for Gravitational Wave Astronomy, UTB. S.B. and K.K. acknowledge support from the German Science Foundation SFB/TR7 “Gravitational Wave Astronomy.” L.C. acknowledges National Science Foundation Grants Number PHY-0653550 and PHY-0955773. LL was supported in part by an NSERC through Discovery Grant. Research at Perimeter Institute is supported through Industry Canada and by the Province of Ontario through the Ministry of Research & Innovation. P.C.D. acknowledges Spanish Ministerio de Educación y Ciencia grant number AYA 2010-21097-C03-01, Generalitat Valenciana grant number PROMETEO-2009-103 and ERC Starting Grant number CAMAP-259276. The work of C.F. is under the auspices of the US Department of Energy, and supported by its contract W-7405-ENG-36 to Los Alamos National Laboratory. L.S.F. acknowledges National Science Foundation Grants Number PHY-0653462, CBET-0940924 and PHY-0969857. B.G. and R.P. acknowledge support from NSF Grant Number AST-1009396 and NASA Grant Number NNX12AO67G. P.L. acknowledges National Science Foundation Grants Number 0903973 and 1205864. D.M.S. acknowledges National Science Foundation Grants Number PHY-0925345 and PHY-0955825. We wish to thank Christian Ott for useful contributions to this review.

-
- [1] J. M. Weisberg and J. H. Taylor. Relativistic Binary Pulsar B1913+16: Thirty Years of Observations and Analysis. *ASP Conf. Ser.*, 328:25, 2005.
 - [2] G. M. Harry and the LIGO Scientific Collaboration. Advanced LIGO: the next generation of gravitational wave detectors. *Classical and Quantum Gravity*, 27(8):084006+, April 2010.
 - [3] Virgo Collaboration. Advanced virgo baseline design. Virgo Technical Report VIR-0027A-09, 2009. <https://tds.ego-gw.it/itf/tds/file.php?callFile=VIR-0027A-09.pdf>.
 - [4] B. Willke, P. Ajith, B. Allen, P. Aufmuth, C. Aulbert, S. Babak, R. Balasubramanian, et al. The GEO-HF project. *Classical and Quantum Gravity*, 23:207, April 2006.
 - [5] G. H. Janssen et al. European Pulsar Timing Array. *AIP Conference Proceedings*, 983(1):633, 2008.
 - [6] F. Jenet et al. The North American Nanohertz Observatory for Gravitational Waves, 2009. arXiv:0909.1058.
 - [7] G. B. Hobbs et al. Gravitational wave detection using pulsars: status of the Parkes Pulsar Timing Array project, 2008. arXiv:0812.2721.
 - [8] B. Iyer et al. LIGO India. Technical report, 2011. <https://dcc.ligo.org/cgi->

- bin/DocDB/ShowDocument?docid=75988.
- [9] K. Somiya. Detector configuration of KAGRA-the Japanese cryogenic gravitational-wave detector. *Classical and Quantum Gravity*, 29(12):124007, June 2012.
 - [10] M. Punturo, M. Abernathy, F. Acernese, B. Allen, N. Andersson, K. Arun, F. Barone, B. Barr, et al. The third generation of gravitational wave observatories and their science reach, April 2010.
 - [11] S. Hild, M. Abernathy, F. Acernese, P. Amaro-Seoane, N. Andersson, K. Arun, F. Barone, B. Barr, et al. Sensitivity studies for third-generation gravitational wave observatories. *Classical and Quantum Gravity*, 28(9):094013, May 2011.
 - [12] K. Danzmann et al. LISA: laser interferometer space antenna for gravitational wave measurements. *Classical and Quantum Gravity*, 13(11A):A247, 1996.
 - [13] P. Amaro-Seoane et al. eLISA: Astrophysics and cosmology in the millihertz regime, 2012. arXiv:1201.3621.
 - [14] B. Abbott et al. LIGO: The Laser Interferometer Gravitational-Wave Observatory. *Rept. Prog. Phys.*, 72:076901, 2009.
 - [15] T. Accadia, F. Acernese, M. Alshourbagy, P. Amico, F. Antonucci, S. Aoudia, N. Arnaud, C. Arnault, K. G. Arun, P. Astone, et al. Virgo: a laser interferometer to detect gravitational waves. *Journal of Instrumentation*, 7:3012, March 2012.
 - [16] H. Grote and LIGO Scientific Collaboration. The GEO 600 status. *Classical and Quantum Gravity*, 27(8):084003, April 2010.
 - [17] Committee for a Decadal Survey of Astronomy and Astrophysics; National Research Council. *New Worlds, New Horizons in Astronomy and Astrophysics*. The National Academies Press, 2010.
 - [18] D. R. Lorimer. Binary and Millisecond Pulsars. *Living Reviews in Relativity*, 11:8, 2008.
 - [19] K. Belczynski, M. Dominik, T. Bulik, R. O’Shaughnessy, C. Fryer, and D. E. Holz. The Effect of Metallicity on the Detection Prospects for Gravitational Waves. *Astrophys. J. Lett.*, 715:L138, 2010.
 - [20] M. Dominik, K. Belczynski, C. Fryer, D. Holz, E. Berti, T. Bulik, I. Mandel, and R. O’Shaughnessy. Double Compact Objects I: The Significance Of The Common Envelope On Merger Rates, February 2012. arXiv:1202.4901.
 - [21] R. Perna and K. Belczynski. Short Gamma-Ray Bursts and Mergers of Compact Objects: Observational Constraints. *Astrophys. J.*, 570:252, 2002.
 - [22] K. Belczynski, R. Perna, T. Bulik, V. Kalogera, N. Ivanova, and D. Q. Lamb. A Study of Compact Object Mergers as Short Gamma-Ray Burst Progenitors. *Astrophys. J.*, 648:1110, 2006.
 - [23] R. O’Shaughnessy, K. Belczynski, and V. Kalogera. Short Gamma-Ray Bursts and Binary Mergers in Spiral and Elliptical Galaxies: Redshift Distribution and Hosts. *Astrophys. J.*, 675:566, 2008.
 - [24] D. Coward, E. Howell, T. Piran, G. Stratta, M. Branchesi, O. Bromberg, B. Gendre, R. Burman, and D. Guetta. The Swift short gamma-ray burst rate density: implications for binary neutron star merger rates, February 2012. arXiv:1202.2179.
 - [25] I. Kowalska, T. Regimbau, T. Bulik, M. Dominik, and K. Belczynski. Effect of metallicity on the gravitational-wave signal from the cosmological population of compact binary coalescences, May 2012. ArXiv:1205.4621.
 - [26] W. Fong, E. Berger, R. Margutti, B. A. Zauderer, E. Troja, I. Czekala, R. Chornock, N. Gehrels, T. Sakamoto, D. B. Fox, and P. Podsiadlowski. A Jet Break in the X-Ray Light Curve of Short GRB 111020A: Implications for Energetics and Rates. *Astrophys. J.*, 756:189, 2012.
 - [27] E. Berger. The Host Galaxies of Short-Duration Gamma-Ray Bursts: Luminosities, Metallicities, and Star-Formation Rates. *Astrophys. J.*, 690:231, 2009.
 - [28] W. Fong, E. Berger, and D. B. Fox. Hubble Space Telescope Observations of Short GRB Host Galaxies: Morphologies, Offsets, and Local Environments. *Astrophys. J.*, 708:9, 2010.
 - [29] E. Troja, S. Rosswog, and N. Gehrels. Precursors of Short Gamma-ray Bursts. *Astrophys. J.*, 723:1711, 2010.
 - [30] R. Margutti, G. Chincarini, J. Granot, C. Guidorzi, E. Berger, M. G. Bernardini, N. Gehrels, A. M. Soderberg, M. Stamatikos, and E. Zaninoni. X-ray flare candidates in short gamma-ray bursts, 2011. arXiv:1107.1740.
 - [31] J. A. Faber and F. A. Rasio. Binary Neutron Star Mergers. *Living Reviews in Relativity*, 15:8, 2012.
 - [32] M. Shibata and K. ō. Uryū. Simulation of merging binary neutron stars in full general relativity: $\Gamma=2$ case. *Phys. Rev. D*, 61(6):064001, 2000.
 - [33] L. Rezzolla, L. Baiotti, B. Giacomazzo, D. Link, and J. A. Font. Accurate evolutions of unequal-mass neutron-star binaries: properties of the torus and short GRB engines. *Class. Quant. Grav.*, 27:114105, 2010.
 - [34] M. Anderson, E. W. Hirschmann, L. Lehner, S. L. Liebling, P. M. Motl, D. Neilsen, C. Palenzuela, and J. E. Tohline. Simulating binary neutron stars: dynamics and gravitational waves. *Phys. Rev.*, D77:024006, 2008.
 - [35] L. Baiotti, B. Giacomazzo, and L. Rezzolla. Accurate evolutions of inspiralling neutron-star binaries: prompt and delayed collapse to black hole. *Phys. Rev. D*, 78:084033, 2008.
 - [36] T. Yamamoto, M. Shibata, and K. Taniguchi. Simulating coalescing compact binaries by a new code SACRA. *Phys. Rev. D*, 78:064054–1–38, 2008.
 - [37] L. Baiotti, B. Giacomazzo, and L. Rezzolla. Accurate evolutions of inspiralling neutron-star binaries: assessment of the truncation error. *Class. Quantum Grav.*, 26:114005, 2009.
 - [38] Y. T. Liu, S. L. Shapiro, Z. B. Etienne, and K. Taniguchi. General relativistic simulations of magnetized binary neutron star mergers. *Phys. Rev. D*, 78:024012, 2008.
 - [39] B. Giacomazzo, L. Rezzolla, and L. Baiotti. Can magnetic fields be detected during the inspiral of binary neutron stars? *MNRAS*, 399:L164, 2009.
 - [40] M. Shibata, K. Taniguchi, and K. Uryū. Merger of binary neutron stars with realistic equations of state in full general relativity. *Phys. Rev. D*, 71(8):084021, 2005.
 - [41] M. Shibata and K. Taniguchi. Merger of binary neutron stars to a black hole: Disk mass, short gamma-ray bursts, and quasinormal mode ringing. *Phys. Rev. D*, 73(6):064027, 2006.
 - [42] E. Berti, V. Cardoso, and A. O. Starinets. TOPICAL REVIEW: Quasinormal modes of black holes and black branes. *Classical and Quantum Gravity*, 26(16):163001, 2009.

- 2009.
- [43] Y. Sekiguchi, K. Kiuchi, K. Kyutoku, and M. Shibata. Effects of Hyperons in Binary Neutron Star Mergers. *Physical Review Letters*, 107(21):211101, 2011.
 - [44] K. Kiuchi, Y. Sekiguchi, K. Kyutoku, and M. Shibata. Gravitational waves, neutrino emissions and effects of hyperons in binary neutron star mergers. *Classical and Quantum Gravity*, 29(12):124003, 2012.
 - [45] K. Kiuchi, Y. Sekiguchi, M. Shibata, and K. Taniguchi. Long-term general relativistic simulation of binary neutron stars collapsing to a black hole. *Phys. Rev. D*, 80(6):064037, 2009.
 - [46] K. Kiuchi, Y. Sekiguchi, M. Shibata, and K. Taniguchi. Exploring Binary-Neutron-Star-Merger Scenario of Short-Gamma-Ray Bursts by Gravitational-Wave Observation. *Physical Review Letters*, 104(14):141101, 2010.
 - [47] Enrico Ramirez-Ruiz. http://online.kitp.ucsb.edu/online/chirps_c12/ramirezruiz/.
 - [48] J. S. Read, C. Markakis, M. Shibata, K. Uryū, J. D. E. Creighton, and J. L. Friedman. Measuring the neutron star equation of state with gravitational wave observations. *Phys. Rev. D*, 79(12):124033, 2009.
 - [49] T. Hinderer, B. D. Lackey, R. N. Lang, and J. S. Read. Tidal deformability of neutron stars with realistic equations of state and their gravitational wave signatures in binary inspiral, 2009. arXiv:0911.3535.
 - [50] C. Markakis, J. S. Read, M. Shibata, K. Uryu, J. Creighton, et al. Neutron star equation of state via gravitational wave observations. *J.Phys.Conf.Ser.*, 189:012024, 2009.
 - [51] F. Özel, G. Baym, and T. Güver. Astrophysical measurement of the equation of state of neutron star matter. *Phys. Rev. D*, 82(10):101301, 2010.
 - [52] F. Özel, A. Gould, and T. Güver. The Mass and Radius of the Neutron Star in the Bulge Low-mass X-Ray Binary KS 1731-260. *Astrophys. J.*, 748:5, 2012.
 - [53] L. Baiotti, T. Damour, B. Giacomazzo, A. Nagar, and L. Rezzolla. Analytic Modeling of Tidal Effects in the Relativistic Inspiral of Binary Neutron Stars. *Physical Review Letters*, 105(26):261101, 2010.
 - [54] L. Baiotti, T. Damour, B. Giacomazzo, A. Nagar, and L. Rezzolla. Accurate numerical simulations of inspiralling binary neutron stars and their comparison with effective-one-body analytical models. *Phys. Rev. D*, 84(2):024017, 2011.
 - [55] S. Bernuzzi, A. Nagar, M. Thierfelder, and B. Brügmann. Tidal effects in binary neutron star coalescence. *Phys. Rev. D*, 86(4):044030, 2012.
 - [56] A. Bauswein and H.-Th. Janka. Measuring neutron-star properties via gravitational waves from binary mergers. *Phys.Rev.Lett.*, 108:011101, 2012.
 - [57] A. Bauswein, H.T. Janka, K. Hebeler, and A. Schwenk. Equation-of-state dependence of the gravitational-wave signal from the ring-down phase of neutron-star mergers. *Phys.Rev.*, D86:063001, 2012.
 - [58] B. Giacomazzo, L. Rezzolla, and L. Baiotti. Accurate evolutions of inspiralling and magnetized neutron stars: Equal-mass binaries. *Phys. Rev. D*, 83(4):044014, 2011.
 - [59] M. Anderson, E. W. Hirschmann, L. Lehner, S. L. Liebling, P. M. Motl, D. Neilsen, C. Palenzuela, and J. E. Tohline. Magnetized Neutron Star Mergers and Gravitational Wave Signals. *Phys. Rev. Lett.*, 100:191101, 2008.
 - [60] L. Rezzolla, B. Giacomazzo, L. Baiotti, J. Granot, C. Kouveliotou, and M. A. Aloy. The Missing Link: Merging Neutron Stars Naturally Produce Jet-like Structures and Can Power Short Gamma-ray Bursts. *Astrophys. J. Lett.*, 732:L6, 2011.
 - [61] L. Lehner, C. Palenzuela, S. L. Liebling, C. Thompson, and C. Hanna. Intense electromagnetic outbursts from collapsing hypermassive neutron stars. *Phys. Rev. D*, 86(10):104035, November 2012.
 - [62] C. Palenzuela, L. Lehner, M. Ponce, S. L. Liebling, M. Anderson, D. Neilsen, and P. Motl. Gravitational and electromagnetic outputs from binary neutron star mergers. *ArXiv e-prints*, January 2013.
 - [63] Y. Sekiguchi, K. Kiuchi, K. Kyutoku, and M. Shibata. Gravitational Waves and Neutrino Emission from the Merger of Binary Neutron Stars. *Physical Review Letters*, 107(5):051102, 2011.
 - [64] R. Gold, S. Bernuzzi, M. Thierfelder, B. Bruegmann, and F. Pretorius. Eccentric binary neutron star mergers, 2011. arXiv:1109.5128.
 - [65] W. E. East, S. T. McWilliams, J. Levin, and F. Pretorius. Observing complete gravitational wave signals from dynamical capture binaries, December 2012. arXiv:1212.0837.
 - [66] W. H. Lee, E. Ramirez-Ruiz, and G. van de Ven. Short Gamma-ray Bursts from Dynamically Assembled Compact Binaries in Globular Clusters: Pathways, Rates, Hydrodynamics, and Cosmological Setting. *Astrophys. J.*, 720:953, 2010.
 - [67] M. Shibata and K. Taniguchi. Coalescence of Black Hole-Neutron Star Binaries. *Living Rev. Relativity*, 14:6, 2011.
 - [68] M. Shibata and K. Uryu. Merger of black hole-neutron star binaries: nonspinning black hole case. *Phys. Rev. D*, 74:121503, 2006.
 - [69] M. Shibata and K. Uryu. Merger of black hole neutron star binaries in full general relativity. *Classical and Quantum Gravity*, 24:125, 2007.
 - [70] M. Shibata and K. Taniguchi. Merger of black hole and neutron star in general relativity: Tidal disruption, torus mass, and gravitational waves. *Phys. Rev. D*, 77(8):084015, 2008.
 - [71] Z. B. Etienne, J. A. Faber, Y. T. Liu, S. L. Shapiro, K. Taniguchi, and T. W. Baumgarte. Fully general relativistic simulations of black hole-neutron star mergers. *Phys. Rev. D*, 77(8):084002, 2008.
 - [72] M. D. Duez, F. Foucart, L. E. Kidder, H. P. Pfeiffer, M. A. Scheel, and S. A. Teukolsky. Evolving black hole-neutron star binaries in general relativity using pseudospectral and finite difference methods. *Phys. Rev. D*, 78(10):104015, 2008.
 - [73] B. Giacomazzo, R. Perna, L. Rezzolla, E. Troja, and D. Lazzati. Compact Binary Progenitors of Short Gamma-Ray Bursts. *Astrophys. J. Lett.*, 762:L18, 2013.
 - [74] Z. B. Etienne, Y. T. Liu, S. L. Shapiro, and T. W. Baumgarte. General relativistic simulations of black-hole-neutron-star mergers: Effects of black-hole spin. *Phys. Rev. D*, 79(4):044024, 2009.
 - [75] F. Foucart, M. D. Duez, L. E. Kidder, M. A. Scheel, B. Szilagyi, and S. A. Teukolsky. Black hole-neutron star mergers for 10MM \odot black holes. *Phys. Rev. D*, 85(4):044015, 2012.
 - [76] S. Chawla, M. Anderson, M. Besselman, L. Lehner, S. L.

- Liebling, P. M. Motl, and D. Neilsen. Mergers of Magnetized Neutron Stars with Spinning Black Holes: Disruption, Accretion, and Fallback. *Physical Review Letters*, 105(11):111101, September 2010.
- [77] F. Pannarale, A. Tonita, and L. Rezzolla. Black Hole-Neutron Star Mergers and Short Gamma-ray Bursts: A Relativistic Toy Model to Estimate the Mass of the Torus. *Astrophys. J.*, 727:95, 2011.
- [78] F. Foucart. Black Hole-Neutron Star Mergers: Disk Mass Predictions, 2012. arXiv:1207.6304.
- [79] Francesco Pannarale. The Black Hole Remnant of Black Hole-Neutron Star Coalescing Binaries, 2012. arxiv:1208.5869.
- [80] M. Shibata, K. Kyutoku, T. Yamamoto, and K. Taniguchi. Gravitational waves from black hole-neutron star binaries: Classification of waveforms. *Phys. Rev. D*, 79(4):044030, 2009.
- [81] F. Pannarale, L. Rezzolla, F. Ohme, and J. S. Read. Will black hole-neutron star binary inspirals tell us about the neutron star equation of state? *Phys. Rev. D*, 84(10):104017, 2011.
- [82] B. D. Lackey, K. Kyutoku, M. Shibata, P. R. Brady, and J. L. Friedman. Extracting equation of state parameters from black hole-neutron star mergers: Nonspinning black holes. *Phys. Rev. D*, 85(4):044061, 2012.
- [83] B. D. Lackey, K. Kyutoku, M. Shibata, P. R. Brady, and J. L. Friedman. Extracting equation of state parameters from black hole-neutron star mergers: aligned-spin black holes and a preliminary waveform model. *ArXiv e-prints*, March 2013.
- [84] M. D. Duez, F. Foucart, L. E. Kidder, C. D. Ott, and S. A. Teukolsky. Equation of state effects in black hole-neutron star mergers. *Classical and Quantum Gravity*, 27(11):114106, 2010.
- [85] K. Kyutoku, M. Shibata, and K. Taniguchi. Gravitational waves from nonspinning black hole-neutron star binaries: Dependence on equations of state. *Phys. Rev. D*, 82(4):044049, 2010.
- [86] K. Kyutoku, H. Okawa, M. Shibata, and K. Taniguchi. Gravitational waves from spinning black hole-neutron star binaries: dependence on black hole spins and on neutron star equations of state. *Phys. Rev. D*, 84(6):064018, 2011.
- [87] F. Foucart, M. D. Duez, L. E. Kidder, and S. A. Teukolsky. Black hole-neutron star mergers: Effects of the orientation of the black hole spin. *Phys. Rev. D*, 83(2):024005, 2011.
- [88] S. Chawla, M. Anderson, M. Besselman, L. Lehner, S. L. Liebling, P. M. Motl, and D. Neilsen. Mergers of Magnetized Neutron Stars with Spinning Black Holes: Disruption, Accretion, and Fallback. *Physical Review Letters*, 105(11):111101, 2010.
- [89] Z. B. Etienne, Y. T. Liu, V. Paschalidis, and S. L. Shapiro. General relativistic simulations of black-hole-neutron-star mergers: Effects of magnetic fields. *Phys. Rev. D*, 85(6):064029, 2012.
- [90] B. M. S. Hansen and M. Lyutikov. Radio and X-ray signatures of merging neutron stars. *Mon. Not. Roy. Astron. Soc.*, 322:695–701, April 2001.
- [91] C. Palenzuela, T. Garrett, L. Lehner, and S. L. Liebling. Magnetospheres of black hole systems in force-free plasma. *Phys. Rev. D*, 82(4):044045, August 2010.
- [92] D. Neilsen, L. Lehner, C. Palenzuela, E. W. Hirschmann, S. L. Liebling, P. M. Motl, and T. Garrett. Boosting jet power in black hole spacetimes. *Proceedings of the National Academy of Science*, 108:12641–12646, August 2011.
- [93] S. T. McWilliams and J. Levin. Electromagnetic Extraction of Energy from Black-hole-Neutron-star Binaries. *Astrophys. J.*, 742:90, December 2011.
- [94] B. C. Stephens, W. E. East, and F. Pretorius. Eccentric Black-hole-Neutron-star Mergers. *Astrophys. J. Lett.*, 737:L5, 2011.
- [95] W. E. East, F. Pretorius, and B. C. Stephens. Eccentric black hole-neutron star mergers: Effects of black hole spin and equation of state. *Phys. Rev. D*, 85(12):124009, 2012.
- [96] I. Mandel and R. O’Shaughnessy. Compact binary coalescences in the band of ground-based gravitational-wave detectors. *Classical and Quantum Gravity*, 27(11):114007, 2010.
- [97] C. Kim, V. Kalogera, and D. R. Lorimer. The Probability Distribution of Binary Pulsar Coalescence Rates. I. Double Neutron Star Systems in the Galactic Field. *Astrophys. J.*, 584:985, 2003.
- [98] R. O’Shaughnessy, V. Kalogera, and K. Belczynski. Mapping Population Synthesis Event Rates on Model Parameters. II. Convergence and Accuracy of Multidimensional Fits. *Astrophys. J.*, 667:1048, 2007.
- [99] K. Belczynski, T. Bulik, I. Mandel, B. S. Sathyaprakash, A. Zdziarski, and J. Mikolajewska. Cyg X-3: a Galactic double black hole or black hole-neutron star progenitor. *ArXiv e-prints*, September 2012.
- [100] M. Dominik, K. Belczynski, C. Fryer, D. E. Holz, E. Berti, T. Bulik, I. Mandel, and R. O’Shaughnessy. Double Compact Objects. I. The Significance of the Common Envelope on Merger Rates. *Astrophys. J.*, 759:52, 2012.
- [101] J. Abadie et al. Predictions for the Rates of Compact Binary Coalescences Observable by Ground-based Gravitational-wave Detectors. *Class.Quant.Grav.*, 27:173001, 2010.
- [102] R. O’Shaughnessy, V. Kalogera, and K. Belczynski. Binary Compact Object Coalescence Rates: The Role of Elliptical Galaxies. *Astrophys. J.*, 716:615, 2010.
- [103] Dimitrios Psaltis. Probes and tests of strong-field gravity with observations in the electromagnetic spectrum. *Living Reviews in Relativity*, 11(9), 2008.
- [104] M. A. Abramowicz, W. Kluzniak, and J. P. Lasota. No observational proof of the black hole event-horizon. *Astron.Astrophys.*, 396:L31, 2002.
- [105] U. Sperhake, E. Berti, and V. Cardoso. Numerical simulations of black-hole binaries and gravitational wave emission, 2011. arXiv:1107.2819.
- [106] T. Bulik, K. Belczynski, and A. Prestwich. IC10 X-1/NGC300 X-1: The Very Immediate Progenitors of BH-BH Binaries. *Astrophys. J.*, 730:140, 2011.
- [107] A. Sadowski, K. Belczynski, T. Bulik, N. Ivanova, F. A. Rasio, and R. O’Shaughnessy. The Total Merger Rate of Compact Object Binaries in the Local Universe. *Astrophys. J.*, 676:1162, 2008.
- [108] M. C. Miller and V. M. Lauburg. Mergers of Stellar-Mass Black Holes in Nuclear Star Clusters. *Astrophys. J.*, 692:917, 2009.
- [109] R. M. O’Leary, B. Kocsis, and A. Loeb. Gravitational waves from scattering of stellar-mass black holes in galactic nuclei. *Mon. Not. Roy. Astron. Soc.*, 395:2127, 2009.

- [110] S. Banerjee, H. Baumgardt, and P. Kroupa. Stellar-mass black holes in star clusters: implications for gravitational wave radiation. *Mon. Not. Roy. Astron. Soc.*, 402:371, 2010.
- [111] F. Pretorius. Evolution of binary black hole spacetimes. *Phys. Rev. Lett.*, 95:121101, 2005.
- [112] M. Campanelli, C. O. Lousto, P. Marronetti, and Y. Zlochower. Accurate evolutions of orbiting black-hole binaries without excision. *Phys. Rev. Lett.*, 96:111101, 2006.
- [113] J. G. Baker, J. Centrella, D.-I. Choi, M. Koppitz, and J. van Meter. Gravitational-Wave Extraction from an Inspiral Configuration of Merging Black Holes. *Physical Review Letters*, 96(11):111102, March 2006.
- [114] Frans Pretorius. Binary Black Hole Coalescence. 2007.
- [115] Harald P. Pfeiffer. Numerical simulations of compact object binaries. *Class.Quant.Grav.*, 29:124004, 2012.
- [116] A. Buonanno et al. Toward faithful templates for non-spinning binary black holes using the effective-one-body approach. *Phys. Rev. D*, 76:104049, 2007.
- [117] P. Ajith et al. Inspiral-merger-ringdown waveforms for black-hole binaries with non-precessing spins. *Phys.Rev.Lett.*, 106:241101, 2011.
- [118] L. Santamaria, F. Ohme, P. Ajith, B. Bruegmann, N. Dorband, et al. Matching post-Newtonian and numerical relativity waveforms: systematic errors and a new phenomenological model for non-precessing black hole binaries. *Phys.Rev.*, D82:064016, 2010.
- [119] R. Sturani et al. Complete phenomenological gravitational waveforms from spinning coalescing binaries. *J.Phys.Conf.Ser.*, 243:012007, 2010.
- [120] A. Taracchini et al. Prototype effective-one-body model for nonprecessing spinning inspiral-merger-ringdown waveforms. *Phys.Rev.*, D86:024011, 2012.
- [121] T. Damour, A. Nagar, and S. Bernuzzi. Improved effective-one-body description of coalescing nonspinning black-hole binaries and its numerical-relativity completion, December 2012. arXiv:1212.4357.
- [122] M. C. Miller and E. J. M. Colbert. Intermediate-Mass Black Holes. *International Journal of Modern Physics D*, 13:1, 2004.
- [123] M. C. Miller. Intermediate-mass black holes as LISA sources. *Classical and Quantum Gravity*, 26(9):094031, 2009.
- [124] D. A. Brown, J. Brink, H. Fang, J. R. Gair, C. Li, G. Lovelace, I. Mandel, and K. S. Thorne. Prospects for Detection of Gravitational Waves from Intermediate-Mass-Ratio Inspirals. *Physical Review Letters*, 99(20):201102, 2007.
- [125] I. Mandel, D. A. Brown, J. R. Gair, and M. C. Miller. Rates and Characteristics of Intermediate Mass Ratio Inspirals Detectable by Advanced LIGO. *Astrophys. J.*, 681:1431, 2008.
- [126] J. M. Fregeau, S. L. Larson, M. C. Miller, R. O’Shaughnessy, and F. A. Rasio. Observing IMBH-IMBH Binary Coalescences via Gravitational Radiation. *Astrophysical Journal Letters*, 646:L135, 2006.
- [127] P. Amaro-Seoane and M. Freitag. Intermediate-Mass Black Holes in Colliding Clusters: Implications for Lower Frequency Gravitational-Wave Astronomy. *Astrophys. J. Lett.*, 653:L53, 2006.
- [128] K. Glampedakis and D. Kennefick. Zoom and whirl: Eccentric equatorial orbits around spinning black holes and their evolution under gravitational radiation reaction. *Phys. Rev. D*, 66(4):044002, 2002.
- [129] J. Levin and G. Perez-Giz. A periodic table for black hole orbits. *Phys. Rev. D*, 77(10):103005, 2008.
- [130] J. Levin and R. Grossman. Dynamics of black hole pairs. I. Periodic tables. *Phys. Rev. D*, 79(4):043016, 2009.
- [131] J. Levin. Energy level diagrams for black hole orbits. *Classical and Quantum Gravity*, 26(23):235010, 2009.
- [132] J. Healy, J. Levin, and D. Shoemaker. Zoom-Whirl Orbits in Black Hole Binaries. *Physical Review Letters*, 103(13):131101, 2009.
- [133] F. Pretorius and D. Khurana. Black hole mergers and unstable circular orbits. *Classical and Quantum Gravity*, 24:83, 2007.
- [134] R. Gold and B. Bruegmann. Radiation from low-momentum zoom-whirl orbits. *Class.Quant.Grav.*, 27:084035, 2010.
- [135] B. Kocsis, M. E. Gáspár, and S. Márka. Detection Rate Estimates of Gravity Waves Emitted during Parabolic Encounters of Stellar Black Holes in Globular Clusters. *Astrophys. J.*, 648:411, 2006.
- [136] R. M. O’Leary, B. Kocsis, and A. Loeb. Gravitational waves from scattering of stellar-mass black holes in galactic nuclei. *Mon. Not. Roy. Astron. Soc.*, 395:2127, 2009.
- [137] S. E. Woosley and J. S. Bloom. The Supernova Gamma-Ray Burst Connection. *Ann. Rev. Astron. Astroph.*, 44:507, 2006.
- [138] J. Hjorth and J. S. Bloom. *The Gamma-Ray Burst - Supernova Connection*, pages 169–190. November 2012.
- [139] A. V. Filippenko. Optical Spectra of Supernovae. *Ann. Rev. Astron. Astroph.*, 35:309, 1997.
- [140] S. R. Kulkarni, D. A. Frail, M. H. Wieringa, R. D. Ekers, E. M. Sadler, R. M. Wark, J. L. Higdon, E. S. Phinney, and J. S. Bloom. Radio emission from the unusual supernova 1998bw and its association with the γ -ray burst of 25 April 1998. *Nature (London)*, 395:663, 1998.
- [141] T. Piran. The physics of gamma-ray bursts. *Reviews of Modern Physics*, 76:1143, October 2004.
- [142] A. M. Soderberg, , et al. Relativistic ejecta from X-ray flash XRF 060218 and the rate of cosmic explosions. *Nature (London)*, 442:1014, 2006.
- [143] A. M. Soderberg et al. A relativistic type Ibc supernova without a detected γ -ray burst. *Nature (London)*, 463:513, 2010.
- [144] S. J. Smartt. Progenitors of Core-Collapse Supernovae. *Ann. Rev. Astron. Astroph.*, 47:63, 2009.
- [145] S. E. Woosley, N. Langer, and T. A. Weaver. The Presupernova Evolution and Explosion of Helium Stars That Experience Mass Loss. *Astrophys. J.*, 448:315, 1995.
- [146] P. Podsiadlowski, P. C. Joss, and J. J. L. Hsu. Presupernova evolution in massive interacting binaries. *Astrophys. J.*, 391:246, 1992.
- [147] S. E. Woosley and A. Heger. The Progenitor Stars of Gamma-Ray Bursts. *Astrophys. J.*, 637:914, 2006.
- [148] S. Taubenberger, S. Valenti, S. Benetti, E. Cappellaro, M. Della Valle, N. Elias-Rosa, S. Hachinger, W. Hillebrandt, K. Maeda, P. A. Mazzali, A. Pastorello, F. Patat, S. A. Sim, and M. Turatto. Nebular emission-line profiles of Type Ib/c supernovae - probing the ejecta asphericity. *Mon. Not. Roy. Astron. Soc.*, 397:677, 2009.
- [149] R. Chornock, A. V. Filippenko, W. Li, and J. M. Silverman. Large Late-Time Asphericities in Three Type IIP Supernovae. *Astrophys. J.*, 713:1363, 2010.

- [150] R. A. Fesen, M. C. Hammell, J. Morse, R. A. Chevalier, K. J. Borkowski, M. A. Dopita, C. L. Gerardy, S. S. Lawrence, J. C. Raymond, and S. van den Bergh. The Expansion Asymmetry and Age of the Cassiopeia A Supernova Remnant. *Astrophys. J.*, 645:283, 2006.
- [151] H. A. Bethe. Supernova mechanisms. *Rev. Mod. Phys.*, 62:801, 1990.
- [152] H.-T. Janka, K. Langanke, A. Marek, G. Martínez-Pinedo, and B. Müller. Theory of core-collapse supernovae. *Phys. Rep.*, 442:38, 2007.
- [153] C. L. Fryer and M. S. Warren. The Collapse of Rotating Massive Stars in Three Dimensions. *Astrophys. J.*, 601:391, 2004.
- [154] A. Marek and H.-T. Janka. Delayed Neutrino-Driven Supernova Explosions Aided by the Standing Accretion-Shock Instability. *Astrophys. J.*, 694:664, 2009.
- [155] K. N. Yakunin, P. Marronetti, A. Mezzacappa, S. W. Bruenn, C.-T. Lee, M. A. Chertkow, W. R. Hix, J. M. Blondin, E. J. Lentz, O. E. Bronson Messer, and S. Yoshida. Gravitational waves from core collapse supernovae. *Class. Quantum Grav.*, 27(19):194005, 2010.
- [156] J. Nordhaus, A. Burrows, A. Almgren, and J. Bell. Dimension as a Key to the Neutrino Mechanism of Core-Collapse Supernova Explosions. *Astrophys. J.*, 720:694, 2010.
- [157] J. W. Murphy and A. Burrows. Criteria for Core-Collapse Supernova Explosions by the Neutrino Mechanism. *Astrophys. J.*, 688:1159, 2008.
- [158] F. Hanke, A. Marek, B. Mueller, and H.-T. Janka. Is Strong SASI Activity the Key to Successful Neutrino-Driven Supernova Explosions? *Submitted to the Astrophys. J.*, *arXiv:1108.4355*, August 2011.
- [159] T. Takiwaki, K. Kotake, and Y. Suwa. Three-dimensional Hydrodynamic Core-Collapse Supernova Simulations for an $11.2 M_{\odot}$ Star with Spectral Neutrino Transport. *Submitted to the Astrophys. J.*, *arXiv:1108.3989*, August 2011.
- [160] A. Burrows, L. Dessart, E. Livne, C. D. Ott, and J. Murphy. Simulations of Magnetically Driven Supernova and Hypernova Explosions in the Context of Rapid Rotation. *Astrophys. J.*, 664:416, 2007.
- [161] S. A. Balbus and J. F. Hawley. A powerful local shear instability in weakly magnetized disks. I - Linear analysis. II - Nonlinear evolution. *Astrophys. J.*, 376:214, 1991.
- [162] M. Obergaulinger, P. Cerdá-Durán, E. Müller, and M. A. Aloy. Semi-global simulations of the magnetorotational instability in core collapse supernovae. *Astron. Astrophys.*, 498:241, 2009.
- [163] P. Cerdá-Durán, J. A. Font, L. Antón, and E. Müller. A new general relativistic magnetohydrodynamics code for dynamical spacetimes. *Astron. Astrophys.*, 492:937, 2008.
- [164] T. Takiwaki and K. Kotake. Gravitational-Wave Signatures in Magnetically-driven Supernova Explosions. *to appear in ApJ*; *arXiv:1004.2896*, April 2010.
- [165] S. M. Couch, J. C. Wheeler, and M. Milosavljević. Aspherical Core-Collapse Supernovae in Red Supergiants Powered by Nonrelativistic Jets. *Astrophys. J.*, 696:953, 2009.
- [166] A. Heger, S.E. Woosley, and H.C. Spruit. Presupernova evolution of differentially rotating massive stars including magnetic fields. *Astrophys. J.*, 626:350, 2005.
- [167] C. D. Ott, A. Burrows, T. A. Thompson, E. Livne, and R. Walder. The Spin Periods and Rotational Profiles of Neutron Stars at Birth. *Astrophys. J. Suppl. Ser.*, 164:130, 2006.
- [168] A. Burrows, E. Livne, L. Dessart, C. D. Ott, and J. Murphy. A New Mechanism for Core-Collapse Supernova Explosions. *Astrophys. J.*, 640:878, 2006.
- [169] A. Burrows, E. Livne, L. Dessart, C. D. Ott, and J. Murphy. Features of the Acoustic Mechanism of Core-Collapse Supernova Explosions. *Astrophys. J.*, 655:416, 2007.
- [170] N. N. Weinberg and E. Quataert. Non-linear saturation of g-modes in proto-neutron stars: quieting the acoustic engine. *Mon. Not. Roy. Astron. Soc.*, 387:L64, 2008.
- [171] S. E. Woosley. Gamma-ray bursts from stellar mass accretion disks around black holes. *Astrophys. J.*, 405:273, 1993.
- [172] E. O'Connor and C. D. Ott. Black Hole Formation in Failing Core-Collapse Supernovae. *Astrophys. J.*, 730:70, 2011.
- [173] S. Nagataki, R. Takahashi, A. Mizuta, and T. Takiwaki. Numerical Study of Gamma-Ray Burst Jet Formation in Collapsars. *Astrophys. J.*, 659:512, 2007.
- [174] S. S. Komissarov, N. Vlahakis, A. Königl, and M. V. Barkov. Magnetic acceleration of ultrarelativistic jets in gamma-ray burst sources. *Mon. Not. Roy. Astron. Soc.*, 394:1182, 2009.
- [175] R. Birkel, M. A. Aloy, H.-T. Janka, and E. Müller. Neutrino pair annihilation near accreting, stellar-mass black holes. *Astron. Astrophys.*, 463:51, 2007.
- [176] S. Harikae, K. Kotake, T. Takiwaki, and Y.-i. Sekiguchi. A General Relativistic Ray-tracing Method for Estimating the Energy and Momentum Deposition by Neutrino Pair Annihilation in Collapsars. *Astrophys. J.*, 720:614, 2010.
- [177] A. I. MacFadyen, S. E. Woosley, and A. Heger. Supernovae, Jets, and Collapsars. *Astrophys. J.*, 550:410, 2001.
- [178] C. C. Lindner, M. Milosavljevic, R. Shen, and P. Kumar. Simulations of Accretion Powered Supernovae in the Progenitors of Gamma Ray Bursts. *Submitted to ApJ*, *arXiv:1108.1415*, August 2011.
- [179] L. Dessart, A. Burrows, E. Livne, and C. D. Ott. The Proto-Neutron Star Phase of the Collapsar Model and the Route to Long-Soft Gamma-Ray Bursts and Hypernovae. *Astrophys. J. Lett.*, 673:L43, 2008.
- [180] T. A. Thompson, P. Chang, and E. Quataert. Magnetar Spin-Down, Hyperenergetic Supernovae, and Gamma-Ray Bursts. *Astrophys. J.*, 611:380, 2004.
- [181] N. Bucciantini, E. Quataert, J. Arons, B. D. Metzger, and T. A. Thompson. Relativistic jets and long-duration gamma-ray bursts from the birth of magnetars. *Mon. Not. Roy. Astron. Soc.*, 383:L25, 2008.
- [182] B. D. Metzger, D. Giannios, T. A. Thompson, N. Bucciantini, and E. Quataert. The protomagnetar model for gamma-ray bursts. *Mon. Not. Roy. Astron. Soc.*, 413:2031, 2011.
- [183] J. Weber. Observation of the Thermal Fluctuations of a Gravitational-Wave Detector. *Phys. Rev. Lett.*, 17:1228, 1966.
- [184] R. Ruffini and J. A. Wheeler. Relativistic Cosmology and Space Platforms. In V. Hardy and H. Moore, editors, *Proceedings of the Conference on Space Physics, ESRO, Paris, France*, page 45, 1971.
- [185] C. Fryer and K. C. B. New. Gravitational waves from

- gravitational collapse. *Liv. Rev. Rel.*, 14:1, 2011.
- [186] C. D. Ott. TOPICAL REVIEW: The gravitational-wave signature of core-collapse supernovae. *Class. Quant. Grav.*, 26(6):063001, 2009.
- [187] K. Kotake, K. Sato, and K. Takahashi. Explosion mechanism, neutrino burst and gravitational wave in core-collapse supernovae. *Rep. Prog. Phys.*, 69:971, 2006.
- [188] A. Marek, H.-T. Janka, and E. Müller. Equation-of-state dependent features in shock-oscillation modulated neutrino and gravitational-wave signals from supernovae. *Astron. Astrophys.*, 496:475, 2009.
- [189] J. Murphy, C. D. Ott, and A. Burrows. Gravitational waves from convection and sasi in core-collapse supernovae. *To be submitted to the Astrophys. J.*, 2009.
- [190] E. Müller, H.-T. Janka, and A. Wongwathanarat. Parametrized 3D models of neutrino-driven supernova explosions: Neutrino emission asymmetries and gravitational-wave signals. *Submitted to Astron. Astrophys. arXiv:1106.6301 [astro-ph]*, June 2011.
- [191] K. Kotake, W. Iwakami Nakano, and N. Ohnishi. Effects of Rotation on Stochasticity of Gravitational Waves in Nonlinear Phase of Core-Collapse Supernovae. *Submitted to ApJ. arXiv:1106.0544 [astro-ph]*, June 2011.
- [192] R. Epstein. The generation of gravitational radiation by escaping supernova neutrinos. *Astrophys. J.*, 223:1037, 1978.
- [193] C. D. Ott, A. Burrows, L. Dessart, and E. Livne. A New Mechanism for Gravitational-Wave Emission in Core-Collapse Supernovae. *Phys. Rev. Lett.*, 96(20):201102, 2006.
- [194] H. Dimmelmeier, C. D. Ott, A. Marek, and H.-T. Janka. Gravitational wave burst signal from core collapse of rotating stars. *Phys. Rev. D*, 78(6):064056, 2008.
- [195] H. Dimmelmeier, C. D. Ott, H.-T. Janka, A. Marek, and E. Müller. Generic Gravitational-Wave Signals from the Collapse of Rotating Stellar Cores. *Phys. Rev. Lett.*, 98(25):251101, 2007.
- [196] K. Kotake, S. Yamada, and K. Sato. Gravitational radiation from axisymmetric rotational core collapse. *Phys. Rev. D*, 68(4):044023, 2003.
- [197] S. Scheidegger, R. Käppeli, S. C. Whitehouse, T. Fischer, and M. Liebendörfer. The influence of model parameters on the prediction of gravitational wave signals from stellar core collapse. *Astron. Astrophys.*, 514:A51, 2010.
- [198] C. D. Ott, H. Dimmelmeier, A. Marek, H.-T. Janka, I. Hawke, B. Zink, and E. Schnetter. 3D Collapse of Rotating Stellar Iron Cores in General Relativity Including Deleptonization and a Nuclear Equation of State. *Phys. Rev. Lett.*, 98:261101, 2007.
- [199] K. Kotake, S. Yamada, K. Sato, K. Sumiyoshi, H. Ono, and H. Suzuki. Gravitational radiation from rotational core collapse: Effects of magnetic fields and realistic equations of state. *Phys. Rev. D*, 69(12):124004, 2004.
- [200] M. Obergaulinger, M. A. Aloy, and E. Müller. Axisymmetric simulations of magneto-rotational core collapse: dynamics and gravitational wave signal. *Astron. Astrophys.*, 450:1107, 2006.
- [201] M. Obergaulinger, M. A. Aloy, H. Dimmelmeier, and E. Müller. Axisymmetric simulations of magnetorotational core collapse: approximate inclusion of general relativistic effects. *Astron. Astrophys.*, 457:209, 2006.
- [202] M. Shibata, Y. T. Liu, S. L. Shapiro, and B. C. Stephens. Magnetorotational collapse of massive stellar cores to neutron stars: Simulations in full general relativity. *Phys. Rev. D*, 74(10):104026, 2006.
- [203] N. Stergioulas. Rotating stars in relativity. *Liv. Rev. Rel.*, 6:3, 2003.
- [204] J. M. Centrella, K. C. B. New, L. L. Lowe, and J. D. Brown. Dynamical Rotational Instability at Low T/W . *Astrophys. J. Lett.*, 550:L193, 2001.
- [205] M. Saijo, T. W. Baumgarte, and S. L. Shapiro. One-armed Spiral Instability in Differentially Rotating Stars. *Astrophys. J.*, 595:352, 2003.
- [206] A. L. Watts, N. Andersson, and D. I. Jones. The Nature of Low $T/|W|$ Dynamical Instabilities in Differentially Rotating Stars. *Astrophys. J. Lett.*, 618:L37, 2005.
- [207] C. D. Ott, S. Ou, J. E. Tohline, and A. Burrows. One-armed Spiral Instability in a Low- $T/|W|$ Postbounce Supernova Core. *Astrophys. J.*, 625:L119, 2005.
- [208] M. Saijo and S. Yoshida. Low $T/|W|$ dynamical instability in differentially rotating stars: diagnosis with canonical angular momentum. *Mon. Not. Roy. Astron. Soc.*, 368:1429, 2006.
- [209] S. Scheidegger, T. Fischer, S. C. Whitehouse, and M. Liebendörfer. Gravitational waves from 3D MHD core collapse simulations. *Astron. Astrophys.*, 490:231, 2008.
- [210] G. Corvino, L. Rezzolla, S. Bernuzzi, R. De Pietri, and B. Giacomazzo. On the shear instability in relativistic neutron stars. *Class. Quantum Grav.*, 27(11):114104, 2010.
- [211] C. D. Ott, C. Reisswig, E. Schnetter, E. O'Connor, U. Sperhake, F. Löffler, P. Diener, E. Abdikamalov, I. Hawke, and A. Burrows. Dynamics and Gravitational Wave Signature of Collapsar Formation. *Phys. Rev. Lett.*, 106(16):161103, 2011.
- [212] K. Kiuchi, M. Shibata, P. J. Montero, and J. A. Font. Gravitational Waves from the Papaloizou-Pringle Instability in Black-Hole-Torus Systems. *Phys. Rev. Lett.*, 106:251102, 2011.
- [213] A. L. Piro and E. Pfahl. Fragmentation of Collapsar Disks and the Production of Gravitational Waves. *Astrophys. J.*, 658:1173, 2007.
- [214] O. Korobkin, E. B. Abdikamalov, E. Schnetter, N. Stergioulas, and B. Zink. Stability of general-relativistic accretion disks. *Phys. Rev. D*, 83(4):043007, 2011.
- [215] A. Corsi and P. Mészáros. Gamma-ray Burst Afterglow Plateaus and Gravitational Waves: Multi-messenger Signature of a Millisecond Magnetar? *Astrophys. J.*, 702:1171, 2009.
- [216] A. L. Piro and C. D. Ott. Supernova Fallback onto Magnetars and Propeller-powered Supernovae. *Astrophys. J.*, 736:108, 2011.
- [217] A. Burrows and J. M. Lattimer. The birth of neutron stars. *Astrophys. J.*, 307:178, 1986.
- [218] A. Burrows and J. M. Lattimer. Convection, Type II supernovae, and the early evolution of neutron stars. *Phys. Rep.*, 163:51, 1988.
- [219] J. A. Pons, S. Reddy, M. Prakash, J. M. Lattimer, and J. A. Miralles. Evolution of Proto-Neutron Stars. *Astrophys. J.*, 513:780, 1999.
- [220] G. F. Burgio, V. Ferrari, L. Gualtieri, and H.-J. Schulze. Oscillations of hot, young neutron stars: Gravitational wave frequencies and damping times. *Phys. Rev. D*, 84(4):044017, 2011.
- [221] P. Gressman, L.-M. Lin, W.-M. Suen, N. Stergioulas, and J. L. Friedman. Nonlinear r-modes in neutron

- stars: Instability of an unstable mode. *Phys. Rev. D*, 66(4):041303, 2002.
- [222] R. Bondarescu, S. A. Teukolsky, and I. Wasserman. Spin evolution of accreting neutron stars: Nonlinear development of the r-mode instability. *Phys. Rev. D*, 76(6):064019, 2007.
- [223] R. Bondarescu, S. A. Teukolsky, and I. Wasserman. Spinning down newborn neutron stars: Nonlinear development of the r-mode instability. *Phys. Rev. D*, 79(10):104003, 2009.
- [224] W. Kastaun, B. Willburger, and K. D. Kokkotas. Saturation amplitude of the f-mode instability. *Phys. Rev. D*, 82(10):104036, 2010.
- [225] N. Andersson and K. D. Kokkotas. The R-Mode Instability in Rotating Neutron Stars. *International Journal of Modern Physics D*, 10:381, 2001.
- [226] C. M. Espinoza, A. G. Lyne, B. W. Stappers, and M. Kramer. A study of 315 glitches in the rotation of 102 pulsars. *Mon. Not. Roy. Astron. Soc.*, 414:1679, 2011.
- [227] B. Abbott et al. Searches for Gravitational Waves from Known Pulsars with Science Run 5 LIGO Data. *Astrophys. J.*, 713:671, 2010.
- [228] B. Abbott et al. Beating the Spin-Down Limit on Gravitational Wave Emission from the Crab Pulsar. *Astrophys. J. Lett.*, 683:L45, 2008.
- [229] J. Abadie et al. Beating the Spin-down Limit on Gravitational Wave Emission from the Vela Pulsar. *Astrophys. J.*, 737:93, 2011.
- [230] A. I. Chugunov and C. J. Horowitz. Breaking stress of neutron star crust. *Mon. Not. Roy. Astron. Soc.*, 407:L54, 2010.
- [231] L. Bildsten. Gravitational Radiation and Rotation of Accreting Neutron Stars. *Astrophys. J. Lett.*, 501:L89, 1998.
- [232] N. Andersson, D. I. Jones, and K. D. Kokkotas. Strange stars as persistent sources of gravitational waves. *Mon. Not. Roy. Astron. Soc.*, 337:1224, 2002.
- [233] B. Abbott et al. Search for gravitational wave radiation associated with the pulsating tail of the SGR 1806-20 hyperflare of 27 December 2004 using LIGO. *Phys. Rev. D*, 76(6):062003, 2007.
- [234] J. Abadie et al. Search for Gravitational Wave Bursts from Six Magnetars. *Astrophys. J. Lett.*, 734:L35, 2011.
- [235] J. Abadie et al. Search for gravitational waves associated with the August 2006 timing glitch of the Vela pulsar. *Phys. Rev. D*, 83(4):042001, 2011.
- [236] J. Abadie et al. First Search for Gravitational Waves from the Youngest Known Neutron Star. *Astrophys. J.*, 722:L504, 2010.
- [237] B. Abbott et al. Search for Gravitational-Wave Bursts from Soft Gamma Repeaters. *Physical Review Letters*, 101(21):211102, 2008.
- [238] M. Punturo et al. The Einstein Telescope: a third-generation gravitational wave observatory. *Classical and Quantum Gravity*, 27(19):194002, 2010.
- [239] K. D. Kokkotas. Gravitational Wave Astronomy (With 2 Figures). In S. Röser, editor, *Reviews in Modern Astronomy*, volume 20 of *Reviews in Modern Astronomy*, page 140, 2008.
- [240] N. Andersson, V. Ferrari, D. I. Jones, K. D. Kokkotas, B. Krishnan, J. S. Read, L. Rezzolla, and B. Zink. Gravitational waves from neutron stars: promises and challenges. *General Relativity and Gravitation*, 43:409, 2011.
- [241] R. Prix, S. Giampanis, and C. Messenger. Search method for long-duration gravitational-wave transients from neutron stars. *Phys. Rev. D*, 84(2):023007, July 2011.
- [242] E. Thrane, S. Kandhasamy, C. D. Ott, W. G. Anderson, N. L. Christensen, M. W. Coughlin, S. Dorsher, S. Giampanis, V. Mandic, A. Mytidis, T. Prestegard, P. Raffai, and B. Whiting. Long gravitational-wave transients and associated detection strategies for a network of terrestrial interferometers. *Phys. Rev. D*, 83(8):083004, April 2011.
- [243] S. Chandrasekhar. The Effect of Gravitational Radiation on the Secular Stability of the Maclaurin Spheroid. *Astrophys. J.*, 161:561, 1970.
- [244] J. L. Friedman and B. F. Schutz. Secular instability of rotating Newtonian stars. *Astrophys. J.*, 222:281, 1978.
- [245] M. Saijo, M. Shibata, T. W. Baumgarte, and S. L. Shapiro. Dynamical Bar Instability in Rotating Stars: Effect of General Relativity. *Astrophys. J.*, 548:919, 2001.
- [246] L. Baiotti, R. de Pietri, G. M. Manca, and L. Rezzolla. Accurate simulations of the dynamical bar-mode instability in full general relativity. *Phys. Rev. D*, 75(4):044023, 2007.
- [247] K. D. Camarda, P. Anninos, P. C. Fragile, and J. A. Font. Dynamical Bar-Mode Instability in Differentially Rotating Magnetized Neutron Stars. *Astrophys. J.*, 707:1610, 2009.
- [248] J. M. Centrella, K. C. B. New, L. L. Lowe, and J. D. Brown. Dynamical Rotational Instability at Low T/W. *Astrophys. J. Lett.*, 550:L193, 2001.
- [249] M. Shibata, S. Karino, and Y. Eriguchi. Dynamical instability of differentially rotating stars. *Mon. Not. Roy. Astron. Soc.*, 334:L27, 2002.
- [250] A. L. Watts, N. Andersson, and D. I. Jones. The Nature of Low T/W—Dynamical Instabilities in Differentially Rotating Stars. *Astrophys. J. Lett.*, 618:L37, 2005.
- [251] P. Cerdá-Durán, V. Quilis, and J. A. Font. AMR simulations of the low T/W—bar-mode instability of neutron stars. *Computer Physics Communications*, 177:288, 2007.
- [252] G. Corvino, L. Rezzolla, S. Bernuzzi, R. De Pietri, and B. Giacomazzo. On the shear instability in relativistic neutron stars. *Classical and Quantum Gravity*, 27(11):114104, 2010.
- [253] J. L. Friedman. Upper Limit on the Frequency of Fast Pulsars. *Physical Review Letters*, 51:718, 1983.
- [254] J. R. Ipser and L. Lindblom. The oscillations of rapidly rotating Newtonian stellar models. II - Dissipative effects. *Astrophys. J.*, 373:213, 1991.
- [255] L. Lindblom and G. Mendell. Does gravitational radiation limit the angular velocities of superfluid neutron stars. *Astrophys. J.*, 444:804, 1995.
- [256] D. Lai and S. L. Shapiro. Gravitational radiation from rapidly rotating nascent neutron stars. *Astrophys. J.*, 442:259, 1995.
- [257] A. Noutsos, D. H. F. M. Schnitzeler, E. F. Keane, M. Kramer, and S. Johnston. Pulsar spin-velocity alignment: kinematic ages, birth periods and braking indices. *Mon. Not. Roy. Astron. Soc.*, 430:2281–2301, April 2013.
- [258] N. Andersson, K. Glampedakis, and B. Haskell. Oscilla-

- tions of dissipative superfluid neutron stars. *Phys. Rev. D*, 79(10):103009, 2009.
- [259] E. Gaertig and K. D. Kokkotas. Oscillations of rapidly rotating relativistic stars. *Phys. Rev. D*, 78(6):064063, 2008.
- [260] Burkhard Zink, Oleg Korobkin, Erik Schnetter, and Nikolaos Stergioulas. Frequency band of the f -mode chandrasekhar-friedman-schutz instability. *Phys. Rev. D*, 81(8):084055, 2010.
- [261] E. Gaertig, K. Glampedakis, K. D. Kokkotas, and B. Zink. The f -mode instability in relativistic neutron stars. *Phys. Rev. Lett.*, 107:101102, 2011.
- [262] N. Andersson. TOPICAL REVIEW: Gravitational waves from instabilities in relativistic stars. *Classical and Quantum Gravity*, 20:105, 2003.
- [263] N. Andersson, K. D. Kokkotas, and N. Stergioulas. On the Relevance of the R-Mode Instability for Accreting Neutron Stars and White Dwarfs. *Astrophys. J.*, 516:307, 1999.
- [264] A. L. Watts, B. Krishnan, L. Bildsten, and B. F. Schutz. Detecting gravitational wave emission from the known accreting neutron stars. *Mon. Not. Roy. Astron. Soc.*, 389:839, 2008.
- [265] L. Rezzolla, F. K. Lamb, and S. L. Shapiro. R-Mode Oscillations in Rotating Magnetic Neutron Stars. *Astrophys. J. Lett.*, 531:L139, 2000.
- [266] W. C. G. Ho, N. Andersson, and B. Haskell. Revealing the Physics of r Modes in Low-Mass X-Ray Binaries. *Physical Review Letters*, 107(10):101101, 2011.
- [267] K. Kokkotas and B. Schmidt. Quasi-Normal Modes of Stars and Black Holes. *Living Reviews in Relativity*, 2:2, 1999.
- [268] N. Andersson and K. D. Kokkotas. Gravitational Waves and Pulsating Stars: What Can We Learn from Future Observations? *Physical Review Letters*, 77:4134, 1996.
- [269] N. Andersson and K. D. Kokkotas. Towards gravitational wave asteroseismology. *Mon. Not. Roy. Astron. Soc.*, 299:1059, 1998.
- [270] O. Benhar, V. Ferrari, and L. Gualtieri. Gravitational wave asteroseismology reexamined. *Phys. Rev. D*, 70(12):124015, 2004.
- [271] E. Gaertig and K. D. Kokkotas. Gravitational wave asteroseismology with fast rotating neutron stars. *Phys. Rev. D*, 83(6):064031, 2011.
- [272] A. Passamonti and N. Andersson. Towards real neutron star seismology: Accounting for elasticity and superfluidity. *ArXiv e-prints*, May 2011.
- [273] G. L. Israel, T. Belloni, L. Stella, Y. Rephaeli, D. E. Gruber, P. Casella, S. Dall’Osso, N. Rea, M. Persic, and R. E. Rothschild. The Discovery of Rapid X-Ray Oscillations in the Tail of the SGR 1806-20 Hyperflare. *Astrophys. J. Lett.*, 628:L53, 2005.
- [274] A. L. Watts and T. E. Strohmayer. Detection with RHESSI of High-Frequency X-Ray Oscillations in the Tail of the 2004 Hyperflare from SGR 1806-20. *Astrophys. J. Lett.*, 637:L117, 2006.
- [275] A. L. Watts and T. E. Strohmayer. Neutron star oscillations and QPOs during magnetar flares. *Advances in Space Research*, 40:1446, 2007.
- [276] V. Hambaryan, R. Neuhauser, and K. D. Kokkotas. Bayesian timing analysis of giant flare of SGR 180620 by RXTE PCA. *Astron. Astrophys.*, 528:A45, 2011.
- [277] S. Mereghetti. The strongest cosmic magnets: soft gamma-ray repeaters and anomalous X-ray pulsars. *Astron. Astrophys. Rev.*, 15:225, 2008.
- [278] L. Samuelsson and N. Andersson. Neutron star asteroseismology. Axial crust oscillations in the Cowling approximation. *Mon. Not. Roy. Astron. Soc.*, 374:256, 2007.
- [279] H. Sotani, K. D. Kokkotas, and N. Stergioulas. Torsional oscillations of relativistic stars with dipole magnetic fields. *Mon. Not. Roy. Astron. Soc.*, 375:261, 2007.
- [280] Y. Levin. On the theory of magnetar QPOs. *Mon. Not. Roy. Astron. Soc.*, 377:159, 2007.
- [281] H. Sotani, K. D. Kokkotas, and N. Stergioulas. Alfvén quasi-periodic oscillations in magnetars. *Mon. Not. Roy. Astron. Soc.*, 385:L5, 2008.
- [282] S. K. Lander and D. I. Jones. Oscillations and instabilities in neutron stars with poloidal magnetic fields. *Mon. Not. Roy. Astron. Soc.*, 412:1730, 2011.
- [283] N. Andersson, K. Glampedakis, and L. Samuelsson. Superfluid signatures in magnetar seismology. *Mon. Not. Roy. Astron. Soc.*, 396:894, 2009.
- [284] M. van Hoven and Y. Levin. Magnetar oscillations - I. Strongly coupled dynamics of the crust and the core. *Mon. Not. Roy. Astron. Soc.*, 410:1036, 2011.
- [285] M. Gabler, P. Cerdá Durán, J. A. Font, E. Müller, and N. Stergioulas. Magneto-elastic oscillations and the damping of crustal shear modes in magnetars. *Mon. Not. Roy. Astron. Soc.*, 410:L37, 2011.
- [286] A. Colaiuda and K. D. Kokkotas. Magnetar oscillations in the presence of a crust. *Mon. Not. Roy. Astron. Soc.*, 414:3014, 2011.
- [287] K. Ioka. Magnetic deformation of magnetars for the giant flares of the soft gamma-ray repeaters. *Mon. Not. Roy. Astron. Soc.*, 327:639, 2001.
- [288] K. Kashiwayama and K. Ioka. Magnetar asteroseismology with long-term gravitational waves. *Phys. Rev. D*, 83(8):081302, 2011.
- [289] A. Corsi and B. J. Owen. Maximum gravitational-wave energy emissible in magnetar flares. *Phys. Rev. D*, 83(10):104014, 2011.
- [290] Y. Levin and M. van Hoven. On the excitation of f -modes and torsional modes by magnetar giant flares. *ArXiv e-prints*, March 2011.
- [291] P. D. Lasky, B. Zink, K. D. Kokkotas, and K. Glampedakis. Hydromagnetic Instabilities in Relativistic Neutron Stars. *Astrophys. J. Lett.*, 735:L20, 2011.
- [292] B. Zink, P. D. Lasky, and K. D. Kokkotas. Are gravitational waves from giant magnetar flares observable? *Phys. Rev. D*, 85(2):024030, January 2012.
- [293] R. Ciolfi, S. K. Lander, G. M. Manca, and L. Rezzolla. Instability-driven Evolution of Poloidal Magnetic Fields in Relativistic Stars. *Astrophys. J. Lett.*, 736:L6, 2011.
- [294] T. Sidery, A. Passamonti, and N. Andersson. The dynamics of pulsar glitches: contrasting phenomenology with numerical evolutions. *Mon. Not. Roy. Astron. Soc.*, 405:1061, 2010.
- [295] C. A. van Eysden and A. Melatos. Gravitational radiation from pulsar glitches. *Classical and Quantum Gravity*, 25(22):225020, 2008.
- [296] É. É. Flanagan and T. Hinderer. Constraining neutron-star tidal Love numbers with gravitational-wave detectors. *Phys. Rev. D*, 77(2):021502, 2008.
- [297] T. Hinderer. Tidal Love Numbers of Neutron Stars.

- Astrophys. J.*, 677:1216, 2008.
- [298] A. J. Penner, N. Andersson, D. I. Jones, L. Samuelsson, and I. Hawke. Crustal failure during binary inspiral. *ArXiv e-prints*, September 2011.
 - [299] D. Tsang, J. S. Read, T. Hinderer, A. L. Piro, and R. Bondarescu. Resonant Shattering of Neutron Star Crusts. *ArXiv e-prints*, October 2011.
 - [300] J. Abadie et al. Sensitivity to Gravitational Waves from Compact Binary Coalescences Achieved during LIGO's Fifth and Virgo's First Science Run, 2010. arXiv:1003.2481.
 - [301] J. Abadie et al. Search for Gravitational Waves from Low Mass Compact Binary Coalescence in LIGO's Sixth Science Run and Virgo's Science Runs 2 and 3. *Phys.Rev.*, D85:082002, 2012.
 - [302] S. Babak et al. Searching for gravitational waves from binary coalescence. *ArXiv e-prints*, August 2012.
 - [303] S. Klimentenko et al. Localization of gravitational wave sources with networks of advanced detectors. *Phys. Rev. D*, 83:102001, 2011.
 - [304] B. F. Schutz. Networks of gravitational wave detectors and three figures of merit. *Classical and Quantum Gravity*, 28(12):125023, 2011.
 - [305] K. Arai et al. Status of Japanese gravitational wave detectors. *Classical and Quantum Gravity*, 26(20):204020, 2009.
 - [306] J. Aasi et al. Prospects for Localization of Gravitational Wave Transients by the Advanced LIGO and Advanced Virgo Observatories, 2013. LIGO-DCC-P1200087, arXiv:1304.0670.
 - [307] S. Fairhurst. Triangulation of gravitational wave sources with a network of detectors. *NJP*, 11:123006, 2009.
 - [308] L. Wen and Y. Chen. Geometrical expression for the angular resolution of a network of gravitational-wave detectors. *Phys. Rev. D*, 81:082001, 2010.
 - [309] S. Klimentenko et al. Localization of gravitational wave sources with networks of advanced detectors. *Phys. Rev. D*, 83(10):102001, 2011.
 - [310] S. Nissanke et al. Localizing Compact Binary Inspirals on the Sky Using Ground-based Gravitational Wave Interferometers. *Astrophys. J.*, 739:99, 2011.
 - [311] J. Veitch et al. Estimating parameters of coalescing compact binaries with proposed advanced detector networks. *Phys. Rev. D*, 85:104045, 2012.
 - [312] LIGO Scientific Collaboration, Virgo Collaboration, J. Abadie, B. P. Abbott, R. Abbott, T. D. Abbott, M. Abernathy, T. Accadia, F. Acernese, C. Adams, and et al. Implementation and testing of the first prompt search for gravitational wave transients with electromagnetic counterparts. *Astron. Astrophys.*, 539:A124, 2012.
 - [313] P. A. Evans, J. K. Fridriksson, N. Gehrels, J. Homan, J. P. Osborne, M. Siegel, A. Beardmore, P. Handbauer, J. Gelbord, J. A. Kennea, and et al. Swift Follow-up Observations of Candidate Gravitational-wave Transient Events. *Astrophys. J. Supp. Ser.*, 203:28, 2012.
 - [314] J. S. Bloom et al. Astro2010 Decadal Survey Whitepaper: Coordinated Science in the Gravitational and Electromagnetic Skies. *ArXiv e-prints*, February 2009. 0902.1527.
 - [315] I. Mandel, L. Z. Kelley, and E. Ramirez-Ruiz. Towards improving the prospects for coordinated gravitational-wave and electromagnetic observations. *ArXiv e-prints*, October 2011.
 - [316] T. Bulik and K. Belczyński. Constraints on the Binary Evolution from Chirp Mass Measurements. *Astrophys. J. Lett.*, 589:L37, 2003.
 - [317] N. Yunes. Gravitational Waves from Compact Binaries as Probes of the Universe. *ArXiv e-prints*, December 2011.
 - [318] D. Gerosa, M. Kesden, E. Berti, R. O'Shaughnessy, and U. Sperhake. Resonant-plane locking and spin alignment in stellar-mass black-hole binaries: a diagnostic of compact-binary formation. *ArXiv e-prints*, February 2013.
 - [319] M. Punturo et al. The third generation of gravitational wave observatories and their science reach. *Classical and Quantum Gravity*, 27(8):084007, 2010.
 - [320] E. Chassande-Mottin et al. Multimessenger astronomy with the Einstein Telescope. *General Relativity and Gravitation*, 43:437, 2011.
 - [321] S. Hild et al. Sensitivity studies for third-generation gravitational wave observatories. *Classical and Quantum Gravity*, 28(9):094013, 2011.
 - [322] B. S. Sathyaprakash, B. F. Schutz, and C. Van Den Broeck. Cosmography with the Einstein Telescope. *Classical and Quantum Gravity*, 27(21):215006, 2010.
 - [323] N. Kawai et al. X-ray All-Sky Monitor on JEM of the Space Station - MAXI (Monitor of All-sky X-ray Image). In M. Matsuoka & N. Kawai, editor, *All-Sky X-Ray Observations in the Next Decade*, page 279, 1997.
 - [324] AGILE. <http://agile.asdc.asi.it/>.
 - [325] FERMI. <http://fermi.gsfc.nasa.gov/>.
 - [326] INTEGRAL. <http://www.rssd.esa.int/index.php?project=INTEGRAL&page=index>.
 - [327] SWIFT. <http://swift.gsfc.nasa.gov/>.
 - [328] CHANDRA. <http://chandra.harvard.edu/>.
 - [329] XMM-Newton. <http://heasarc.gsfc.nasa.gov/docs/xmm/xmmgof.html>.
 - [330] SUZAKU. <http://www.astro.isas.jaxa.jp/suzaku/index.html.en/>.
 - [331] NUSTAR. <http://www.nustar.caltech.edu/>.
 - [332] MAXI. <http://maxi.riken.jp>.
 - [333] P. C. Agrawal. A broad spectral band Indian Astronomy satellite *Astrosat*. *Advances in Space Research*, 38:2989, 2006.
 - [334] V. Koteswara Rao et al. The scientific objectives of the ASTROSAT mission of ISRO. *Acta Astronautica*, 65:6, 2009.
 - [335] ASTROSAT. <http://astrosat.iucaa.in>.
 - [336] T. Takahashi et al. The ASTRO-H X-ray Observatory, 2012. arXiv:1210.4378.
 - [337] N. Kawai and D. Yonetoku. Observing Gamma-Ray Bursts with ASTRO-H. In N. Kawai and S. Nagataki, editors, *American Institute of Physics Conference Series*, volume 1279 of *American Institute of Physics Conference Series*, page 204, October 2010.
 - [338] J. Paul et al. The Chinese–French SVOM mission for gamma-ray burst studies. *Comptes Rendus Physique*, 12:298, 2011.
 - [339] S. Schanne et al. The future Gamma-Ray Burst Mission SVOM. *ArXiv e-prints*, May 2010.
 - [340] SVOM. <http://smc.cnes.fr/SVOM/index.htm>.
 - [341] Concepts for Future High-Energy Astrophysics Missions. <http://heasarc.gsfc.nasa.gov/docs/heasarc/missions/concepts.html>.
 - [342] R. Petre et al. The NASA X-ray Mission concepts study. In *Society of Photo-Optical Instrumentation Engineers*

- (SPIE) Conference Series, volume 8443 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, September 2012.
- [343] J. Camp, personal communication.
 - [344] J. Sylvestre. Prospects for the Detection of Electromagnetic Counterparts to Gravitational Wave Events. *Astrophys. J.*, 591:1152, 2003.
 - [345] C. W. Stubbs. Linking optical and infrared observations with gravitational wave sources through transient variability. *Classical and Quantum Gravity*, 25(18):184033, 2008.
 - [346] J. S. Bloom et al. Astro2010 Decadal Survey Whitepaper: Coordinated Science in the Gravitational and Electromagnetic Skies, February 2009. arXiv:0902.1527.
 - [347] E. S. Phinney. Finding and Using Electromagnetic Counterparts of Gravitational Wave Sources. In *astro2010: The Astronomy and Astrophysics Decadal Survey*, volume 2010 of *Astronomy*, page 235, 2009.
 - [348] S. Kulkarni and M. M. Kasliwal. Transients in the Local Universe. In N. Kawai, T. Mihara, M. Kohama, & M. Suzuki, editor, *Astrophysics with All-Sky X-Ray Observations*, page 312, March 2009.
 - [349] LOFAR. <http://www.lofar.org/>.
 - [350] EVLA. <http://www.nrao.edu/index.php/about/facilities/vlaevla>.
 - [351] ASKAP. <http://www.atnf.csiro.au/projects/askap/index.html>.
 - [352] SKA. <http://www.skatelescope.org/>.
 - [353] M. V. Medvedev and A. Loeb. On Poynting-flux-driven bubbles and shocks around merging neutron star binaries. *Mon. Not. Roy. Astron. Soc.*, March 2013.
 - [354] E. Nakar and T. Piran. Detectable radio flares following gravitational waves from mergers of binary neutron stars. *Nature (London)*, 478:82, 2011.
 - [355] B. D. Metzger and E. Berger. What is the Most Promising Electromagnetic Counterpart of a Neutron Star Binary Merger? *Astrophys. J.*, 746:48, 2012.
 - [356] T. Piran, E. Nakar, and S. Rosswog. The electromagnetic signals of compact binary mergers. *Mon. Not. Roy. Astron. Soc.*, page 771, 2013.
 - [357] I. Bartos, P. Brady, and S. Marka. How Gravitational-wave Observations Can Shape the Gamma-ray Burst Paradigm. *ArXiv e-prints*, December 2012.
 - [358] S. Nissanke, M. Kasliwal, and A. Georgieva. Identifying Elusive Electromagnetic Counterparts to Gravitational Wave Mergers: an end-to-end simulation. *ArXiv e-prints*, October 2012.
 - [359] L. Z. Kelley, I. Mandel, and E. Ramirez-Ruiz. Electromagnetic transients as triggers in searches for gravitational waves from compact binary mergers. *ArXiv e-prints*, September 2012.
 - [360] D. Kasen, N. R. Badnell, and J. Barnes. Opacities and Spectra of the r-process Ejecta from Neutron Star Mergers. *ArXiv e-prints*, March 2013.
 - [361] R. M. Bionta et al. Observation of a neutrino burst in coincidence with supernova 1987A in the Large Magellanic Cloud. *Phys. Rev. Lett.*, 58:1494, 1987.
 - [362] K. Hirata et al. Observation of a neutrino burst from the supernova SN1987A. *Phys. Rev. Lett.*, 58:1490, 1987.
 - [363] E.N. Alexeyev, L.N. Alexeyeva, V.I. Volchenko, and I.V. Krivosheina. Possible detection of a neutrino signal on 23 February 1987 at the Baksan underground scintillation telescope of the Institute of Nuclear Research. *JETP Lett.*, 45:589, 1987.
 - [364] M. Aglietta, G. Badino, G. Bologna, C. Castagnoli, A. Castellina, et al. On the event observed in the Mont Blanc Underground Neutrino observatory during the occurrence of Supernova 1987a. *Europhys.Lett.*, 3:1315, 1987.
 - [365] J.N. Bahcall. *Neutrino Astrophysics*. Cambridge University Press, 1989.
 - [366] G. Pagliaroli, F. Vissani, E. Coccia, and W. Fulgione. Neutrinos from Supernovae as a Trigger for Gravitational Wave Search. *Phys. Rev. Lett.*, 103:031102, 2009.
 - [367] Kate Scholberg. Supernova Neutrino Detection. *Ann.Rev.Nucl.Part.Sci.*, 62:81, 2012.
 - [368] W. Fulgione. Status of supernova neutrino detectors. *J. Phys.: Conf. Ser.*, 203:012077, 2010.
 - [369] S. Fukuda et al. The super-kamiokande detector. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 501(2-3):418, 2003.
 - [370] M. Ikeda et al. Search for Supernova Neutrino Bursts at Super-Kamiokande. *The Astrophysical Journal*, 669:519, 2007.
 - [371] W. for the LVD Collaboration Fulgione. The LVD Neutrino Observatory. *Memorie della Societa Astronomica Italiana Supplement*, 9:388, 2006.
 - [372] N. Yu. Agafonova et al. On-line recognition of supernova neutrino bursts in the LVD detector. *Astropart. Phys.*, 28:516, 2008.
 - [373] G. Alimonti et al. The Borexino detector at the Laboratori Nazionali del Gran Sasso. *Nuclear Instruments and Methods in Physics Research A*, 600:568, 2009.
 - [374] L. Cadonati, F. P. Calaprice, and M. C. Chen. Supernova neutrino detection in Borexino. *Astropart. Phys.*, 16:361, 2002.
 - [375] Kamland. <http://www.awa.tohoku.ac.jp/KamLAND/>.
 - [376] A. Piepke. KamLAND: A reactor neutrino experiment testing the solar neutrino anomaly. *Nucl.Phys.Proc.Suppl.*, 91:99, 2001.
 - [377] Sno+. <http://snoplus.phy.queensu.ca/Home.html>.
 - [378] J Ahrens et al. Sensitivity of the icecube detector to astrophysical sources of high energy muon neutrinos. *Astroparticle Physics*, 20(5):507, 2004.
 - [379] F. Halzen, J. E. Jacobsen, and E. Zas. Ultra-Transparent Antarctic Ice as a Supernova Detector. *Phys. Rev.*, D53:7359, 1996.
 - [380] P. Antonioli et al. SNEWS: The SuperNova Early Warning System. *New J. Phys.*, 6:114, 2004.
 - [381] K. Scholberg. The SuperNova Early Warning System. *Astron. Nachr.*, 329:337, 2008.
 - [382] P. J. Sutton. A rule of thumb for the detectability of gravitational-wave bursts. Technical Report LIGO-P1000041-v1, LIGO, 2010. <https://dcc.ligo.org/cgi-bin/private/DocDB/ShowDocument?docid=10614>.
 - [383] C. D. Ott. The Gravitational Wave Signature of Core-Collapse Supernovae. *Class. Quantum Grav.*, 26:063001, 2009. arXiv:0809.0695.
 - [384] W. Fulgione. Status of supernova neutrino detectors. *Journal of Physics Conference Series*, 203(1):012077, 2010.
 - [385] C. D. Ott. Probing the core-collapse supernova mechanism with gravitational waves. *Classical and Quantum Gravity*, 26(20):204015, 2009.
 - [386] N. Arnaud et al. Detection of a close supernova gravitational wave burst in a network of interferometers, neutrino and optical detectors. *Astroparticle Physics*,

- 21:201, 2004.
- [387] I. Leonor et al. Searching for prompt signatures of nearby core-collapse supernovae by a joint analysis of neutrino and gravitational-wave data. *Class. Quant. Grav.*, 27:084019, 2010.
 - [388] G. Pagliaroli et al. Neutrinos from Supernovae as a Trigger for Gravitational Wave Search. *Physical Review Letters*, 103(3):031102, 2009.
 - [389] C. Spiering. Status and Perspectives of Astroparticle Physics in Europe, April 2008. arXiv: 0804.1500.
 - [390] D. Autiero et al. Large underground, liquid based detectors for astro-particle physics in europe: scientific case and prospects. *JCAP*, 0711:011, 2007.
 - [391] M. D. Kistler et al. Core-Collapse Astrophysics with a Five-Megaton Neutrino Detector, 2008. arXiv:0810.1959.
 - [392] S. Ando, J. F. Beacom, and H. Yüksel. Detection of neutrinos from supernovae in nearby galaxies. *Phys. Rev. Lett.*, 95(17):171101, 2005.
 - [393] A. de Bellefon et al. MEMPHYS: A large scale water Cherenkov detector at Fréjus, 2006. arxiv: hep-ex/0607026.
 - [394] M. Wurm et al. The next-generation liquid-scintillator neutrino observatory LENA. *ArXiv e-prints*, April 2011.
 - [395] A. Rubbia. Underground neutrino detectors for particle and astroparticle science: The giant liquid argon charge imaging experiment (glacier). *Journal of Physics: Conference Series*, 171(1):012020, 2009.
 - [396] J. Maricic and the Lbne Dusel collaboration. The Long Baseline Neutrino Oscillation Experiment at DUSEL. *Journal of Physics Conference Series*, 203(1):012109, 2010.
 - [397] K. Nakamura. Hyper-Kamiokande - a Next Generation Water Cherenkov Detector. In R. Shrock, editor, *Neutrinos and Implications for Physics Beyond the Standard Model*, page 307. World Scientific, 2003.
 - [398] Y. Suzuki. Multi-Megaton Water Cherenkov Detector for a Proton Decay Search – TITAND, October 2001. arXiv:hep-ex/0110005.
 - [399] S. Böser, M. Kowalski, L. Schulte, N. Linn Strotjohann, and M. Voge. Detecting extra-galactic supernova neutrinos in the Antarctic ice. *ArXiv e-prints*, April 2013.
 - [400] I. Bartos, B. Dasgupta, and S. Márka. Probing the structure of jet-driven core-collapse supernova and long gamma-ray burst progenitors with high-energy neutrinos. *Phys. Rev. D*, 86(8):083007, 2012.
 - [401] R. Abbasi et al. The design and performance of IceCube DeepCore. *Astroparticle Physics*, 35:615, 2012.
 - [402] I. Bartos et al. Detection Prospects for GeV Neutrinos from Collisionally Heated Gamma-ray Bursts with IceCube/DeepCore. *ArXiv e-prints*, January 2013.
 - [403] M. Ageron et al. ANTARES: The first undersea neutrino telescope. *Nuclear Instruments and Methods in Physics Research A*, 656:11, 2011.
 - [404] M. de Jong. The KM3NeT neutrino telescope. *Nucl. Instrum. Methods Phys. Res., Sect. A*, 623(1):445, 2010.
 - [405] A. Avrorin et al. The baikal neutrino experiment. *Nucl. Instrum. Methods Phys. Res., Sect. A*, 626-627(Supplement 1):S13, 2011.
 - [406] E. Waxman and J. Bahcall. High energy neutrinos from cosmological gamma-ray burst fireballs. *Physical Review Letters*, 78:2292, 1997.
 - [407] D. Guetta et al. Neutrinos from individual gamma-ray bursts in the BATSE catalog. *Astropart. Phys.*, 20(4):429, 2004.
 - [408] J. Ahrens et al. Sensitivity of the IceCube detector to astrophysical sources of high energy muon neutrinos. *Astroparticle Physics*, 20:507, 2004.
 - [409] R. Abbasi et al. An absence of neutrinos associated with cosmic-ray acceleration in γ -ray bursts. *Nature (London)*, 484:351, 2012.
 - [410] S. Hümmer, P. Baerwald, and W. Winter. Neutrino emission from gamma-ray burst fireballs, revised. *Phys. Rev. Lett.*, 108:231101, 2012.
 - [411] H.-N. He et al. Icecube Nondetection of Gamma-Ray Bursts: Constraints on the Fireball Properties. *Astrophys. J.*, 752:29, 2012.
 - [412] S. Ando and J. F. Beacom. Revealing the Supernova–Gamma-Ray Burst Connection with TeV Neutrinos. *Phys. Rev. Lett.*, 95(6):061103, 2005.
 - [413] S. Horiuchi and S. Ando. High-energy neutrinos from reverse shocks in choked and successful relativistic jets. *Phys. Rev. D*, 77(6):063007, 2008.
 - [414] Soebur Razzaque, Peter Mészáros, and Eli Waxman. TeV Neutrinos from Core Collapse Supernovae and Hypernovae. *Phys. Rev. Lett.*, 93(18):181101, 2004. Erratum: *Phys. Rev. Lett.* 94, 109903 (2005).
 - [415] I. Bartos et al. Observational constraints on multimessenger sources of gravitational waves and high-energy neutrinos. *Phys. Rev. Lett.*, 107:251101, 2011.
 - [416] S. Adrian-Martinez et al. A First Search for coincident Gravitational Waves and High Energy Neutrinos using LIGO, Virgo and ANTARES data from 2007, 2012. arXiv:1205.3018.
 - [417] B. Abbott and for the LIGO Scientific Collaboration. Implications for the Origin of GRB 070201 from LIGO Observations. *Astrophys. J.*, 681:1419, 2008.
 - [418] J. Abadie et al. Implications For The Origin Of GRB 051103 From LIGO Observations. *Astrophys. J.*, 755:2, 2012.
 - [419] J. Abadie et al. Search for Gravitational Wave Bursts from Six Magnetars. *Astrophys. J.*, 734:L35, 2011.
 - [420] J. Abadie et al. A search for gravitational waves associated with the August 2006 timing glitch of the Vela pulsar. *Phys.Rev.*, D83:042001, 2011.
 - [421] M.S. Briggs et al. Search for gravitational waves associated with gamma-ray bursts during LIGO science run 6 and Virgo science runs 2 and 3. *Astrophys.J.*, 760:12, 2012.
 - [422] J. Aasi et al. Search for Gravitational Waves from Binary Black Hole Inspiral, Merger and Ringdown in LIGO-Virgo Data from 2009-2010. *Phys. Rev. D*, 87:022002, 2013.
 - [423] J. Abadie et al. Search for Gravitational Waves from Intermediate Mass Binary Black Holes. *Phys.Rev.*, D85:102004, 2012.
 - [424] J. Abadie et al. All-sky search for gravitational-wave bursts in the second joint LIGO-Virgo run. *Phys.Rev.*, D85:122007, 2012.
 - [425] B.P. Abbott et al. Search for gravitational wave ring-downs from perturbed black holes in LIGO S4 data. *Phys.Rev.*, D80:062001, 2009.
 - [426] B.P. Abbott et al. First LIGO search for gravitational wave bursts from cosmic (super)strings. *Phys.Rev.*, D80:062002, 2009.
 - [427] J. Abadie, B. P. Abbott, R. Abbott, M. Abernathy, C. Adams, R. Adhikari, P. Ajith, B. Allen, G. Allen, E. Amador Ceron, and et al. Calibration of the LIGO

- gravitational wave detectors in the fifth science run. *Nuclear Instruments and Methods in Physics Research A*, 624:223–240, December 2010.
- [428] N. Christensen, LIGO Scientific Collaboration, and Virgo Collaboration. LIGO S6 detector characterization studies. *Classical and Quantum Gravity*, 27(19):194010, October 2010.
- [429] M. W. E. Smith, D. B. Fox, D. F. Cowen, P. Mészáros, G. Tešić, J. Fixelle, I. Bartos, P. Sommers, A. Ashtekar, G. Jogesh Babu, S. D. Barthelmy, S. Coutu, T. DeYoung, A. D. Falcone, L. S. Finn, S. Gao, B. Hashemi, A. Homeier, S. Márka, B. J. Owen, and I. Taboada. The Astrophysical Multimessenger Observatory Network (AMON). *Astropart. Phys.*, 45:56, May 2013.
- [430] N. Yunes and X. Siemens. Gravitational Wave Tests of General Relativity with Ground-Based Detectors and Pulsar Timing Arrays. *ArXiv e-prints*, April 2013.
- [431] L. Z. Kelley, E. Ramirez-Ruiz, M. Zemp, J. Diemand, and I. Mandel. The Distribution of Coalescing Compact Binaries in the Local Universe: Prospects for Gravitational-wave Observations. *Astrophys. J. Lett.*, 725:L91–L96, December 2010.
- [432] E. Berger. The environments of short-duration gamma-ray bursts and implications for their progenitors. *New Astron. Rev.*, page 1, 2011.
- [433] B. Aylott et al. Testing gravitational-wave searches with numerical relativity waveforms: Results from the first Numerical INjection Analysis (NINJA) project. *Class. Quant. Grav.*, 26:165008, 2009.
- [434] L. Cadonati et al. Status of NINJA: the Numerical INjection Analysis project. *Classical and Quantum Gravity*, 26(11):114008, 2009.
- [435] The Omega Pipeline. <https://geco.phys.columbia.edu/omega>.
- [436] S. Fischetti, J. Healy, L. Cadonati, L. London, S. R. P. Mohapatra, and D. Shoemaker. Exploring the use of numerical relativity waveforms in burst analysis of precessing black hole mergers. *Phys. Rev. D*, 83(4):044019, 2011.
- [437] A. Heger, N. Langer, and S.E. Woosley. Presupernova Evolution of Rotating Massive Stars. I. Numerical Method and Evolution of the Internal Stellar Structure. *Astrophys. J.*, 528:368, 2000.
- [438] R. Hirschi, G. Meynet, and A. Maeder. Stellar evolution with rotation. xii. pre-supernova models. *Astron. Astroph.*, 443:649, 2005.
- [439] R. Hirschi, G. Meynet, and A. Maeder. Stellar evolution with rotation. xiii. predicted grb rates at various z. *Astron. Astroph.*, 443:581, 2005.
- [440] S.-C. Yoon and N. Langer. Evolution of rapidly rotating metal-poor massive stars towards gamma-ray bursts. *Astron. Astroph.*, 443:643, 2005.
- [441] S. E. Woosley and A. Heger. The Progenitor Stars of Gamma-Ray Bursts. *Astrophys. J.*, 637:914, 2006.
- [442] S.-C. Yoon, N. Langer, and C. Norman. On the evolution of rapidly rotating massive white dwarfs towards supernovae or collapses. *Astron. Astroph.*, 460:199, 2006.
- [443] C.L. Fryer, S.E. Woosley, and D.H. Hartmann. Formation rates of black hole accretion disk gamma-ray bursts. *Astrophys. J.*, 526:152, 1999.
- [444] Ph. Podsiadlowski, P.A. Mazzali, K. Nomoto, D. Lazdazi, and E. Cappellaro. The rates of hypernovae and gamma-ray bursts: Implications for their progenitors. *Astrophys. J.*, 607:L17, 2004.
- [445] C.L. Fryer and A. Heger. Binary Merger Progenitors for Gamma-Ray Bursts and Hypernovae. *Astrophys. J.*, 623:302, 2005.
- [446] E.P.J. van den Heuvel and S.-C. Yoon. Long gamma-ray burst progenitors: boundary conditions and binary models. *Astrophysics and Space Science*, 311:177, 2007.
- [447] M. Cantiello, S.-C. Yoon, N. Langer, and M. Livio. Binary star progenitors of long gamma-ray bursts. *Astron. Astroph.*, 465:L29, 2007.
- [448] S.-C. Yoon, N. Langer, and S. Scheithauer. Effects of rotation on the helium burning shell source in accreting white dwarfs. *Astron. Astroph.*, 425:217, 2004.
- [449] S.-C. Yoon and N. Langer. On the evolution of rapidly rotating massive white dwarfs towards supernovae or collapses. *Astron. Astroph.*, 435:967, 2005.
- [450] S.-C. Yoon, P. Podsiadlowski, and S. & Rosswog. Remnant evolution after a carbon-oxygen white dwarf merger. *Mon. Not. R. Astron. Soc.*, 380:933, 2007.
- [451] C.L. et al. Fryer. Spectra and Light Curves of Failed Supernovae. *Astrophys. J.*, 707:193, 2009.
- [452] L. Dessart, A. Burrows, C. D. Ott, E. Livne, S.-Y. Yoon, and N. Langer. Multidimensional Simulations of the Accretion-induced Collapse of White Dwarfs to Neutron Stars. *Astrophys. J.*, 644:1063, 2006.
- [453] E. B. Abdikamalov, C. D. Ott, L. Rezzolla, L. Dessart, H. Dimmelfmeier, A. Marek, and H.-T. Janka. Axisymmetric general relativistic simulations of the accretion-induced collapse of white dwarfs. *Phys. Rev. D*, 81:044012, 2010.
- [454] G. Dimonte et al. A comparative study of the turbulent Rayleigh-Taylor instability using high-resolution three-dimensional numerical simulations: The Alpha-Group collaboration. *Astrophys. J.*, 16:1668, 2004.
- [455] D.H. Porter and P.R. Woodward. Using ppm to model turbulent stellar convection. In F. Grinstein, L. Margolin, and W. Rider, editors, *Implicit Large Eddy Simulation: Computing Turbulent Fluid Dynamics*, Los Alamos, NM, 2006. Cambridge University Press.
- [456] N. Andersson and G. L. Comer. Probing Neutron-Star Superfluidity with Gravitational-Wave Data. *Physical Review Letters*, 87(24):241101, 2001.
- [457] S. T. McWilliams, J. P. Ostriker, and F. Pretorius. The imminent detection of gravitational waves from massive black-hole binaries with pulsar timing arrays, 2012. arXiv:1211.4590.
- [458] A. Sesana. Systematic investigation of the expected gravitational wave signal from supermassive black hole binaries in the pulsar timing band, 2012. arXiv:1211.5375.